

Iterated algebraic injectivity and the faithfulness conjecture

John Bourke^a

^aDepartment of Mathematics and Statistics, Masaryk University,
Kotlářská 2, Brno 61137, Czech Republic

Abstract

Algebraic injectivity was introduced to capture homotopical structures like algebraic Kan complexes. But at a much simpler level, it allows one to describe sets with operations subject to no equations. If one wishes to add equations (or operations of greater complexity) then it is natural to consider iterated algebraic injectives, which we introduce and study in the present paper.

Our main application concerns Grothendieck’s weak ω -groupoids, introduced in Pursuing Stacks, and the closely related definition of weak ω -category due to Maltsiniotis. Using ω iterations we describe these as iterated algebraic injectives and, via this correspondence, prove the faithfulness conjecture of Maltsiniotis. Through work of Ara, this implies a tight correspondence between the weak ω -categories of Maltsiniotis and those of Batanin/Leinster.

Communicated by: Clemens Berger.

Received: 22nd February, 2019. Accepted: 4th May, 2020.

MSC: 18D05, 18C10, 55U35.

Keywords: algebraic injective, globular theory, faithfulness conjecture.

1. Introduction

The categorical notion of injectivity captures interesting structures in many areas — for instance, injective modules in algebra and Kan complexes in homotopy theory. *Algebraic injectivity* was introduced [13, 20] with the homotopical examples in mind, but even in the simplest settings it has much to say.

For instance, whilst all sets are injective with respect to monomorphisms, an algebraic injective with respect to the inclusion $j : 2 \hookrightarrow 3$ is a set X equipped with a binary operation m , or magma, as depicted below.

$$\begin{array}{ccc} 2 & \xrightarrow{(a,b)} & X \\ j \downarrow & \nearrow & \\ 3 & & \end{array} \quad \begin{array}{c} \\ \\ (a,b,m(a,b)) \end{array}$$

Email address: bourkej@math.muni.cz (John Bourke)

© John Bourke, 2020, under a Creative Commons Attribution 4.0 International License.

DOI: 10.21136/HS.2020.13

Now iterating injectivity is a futile affair: injectives in injectives are injectives. On the other hand, in the present paper we will show that *iterating algebraic injectivity* produces interesting structures. For a simple example, all universal algebraic structures are algebraic injectives in algebraic injectives (a.k.a. *2-injectives*) in **Set** — see Example 3.11 below.

The primary case of interest here concerns the cellular globular theories of [19] used to define Grothendieck weak ω -groupoids [14]. In this setting, we will show that models of cellular globular theories can be identified as certain ω -injectives — iterated algebraic injectives using ω iterations. We will use this identification to prove the *faithfulness conjecture* of Maltsiniotis [19], which we now describe in a little more detail.

Grothendieck’s weak ω -groupoids [14] and the weak ω -categories introduced by Maltsiniotis in [19] are globular sets with additional structure, expressed in terms of their being models of certain *globular theories*. In order that the models of a globular theory \mathbb{T} are equipped with the operations expected in a higher categorical structure one imposes contractibility conditions upon \mathbb{T} . To capture weakness of these operations one requires that \mathbb{T} is *cellular* — this means that it can be constructed as a colimit of a chain

$$\Theta_0 = \mathbb{T}_0 \longrightarrow \mathbb{T}_1 \longrightarrow \dots \longrightarrow \mathbb{T}_n \xrightarrow{J_{n+1}^n} \mathbb{T}_{n+1} \xrightarrow{J_{\omega}^{n+1}} \mathbb{T}$$

in which Θ_0 is the initial globular theory, and in which each $J_{n+1}^n : \mathbb{T}_n \rightarrow \mathbb{T}_{n+1}$ is obtained by *freely adjoining fillers* — see Section 2.4. In [19] Maltsiniotis conjectured that each functor $J_m^n : \mathbb{T}_n \rightarrow \mathbb{T}_m$ for $n < m$ defining a cellular globular theory is faithful. Assuming this conjecture Ara [2] established a sharp correspondence between the weak ω -categories introduced by Maltsiniotis and those of Batanin/Leinster [4, 18]. However the faithfulness conjecture, on which the correspondence depends, was left unproven. Using the framework of iterated algebraic injectivity, we prove it in Theorem 5.2.

Let us now describe the structure of the paper. In Section 2 we describe some background on globular sets, globular theories and the faithfulness conjecture. In Section 3 we recall algebraic injectivity, introduce iterated algebraic injectivity and the important class of $(\mathcal{A}, \mathcal{B})$ -iterated algebraic injectives, which are parametrised by a set of objects \mathcal{A} and family of maps \mathcal{B} in the base category. In Theorem 3.12 we show that, for a suitable choice of \mathcal{A} and \mathcal{B} , these capture the models of cellular globular theories. In Section 4.1 we abstract results of Nikolaus [20] relating to the construction of free algebraic injectives, and extend them to our iterated setting. Using these results, we prove the faithfulness conjecture in Section 5. In Section 6, using the framework of [11], we generalise some of the results in Section 3 away from the globular setting — in particular, we show that in general $(\mathcal{A}, \mathcal{B})$ -iterated algebraic injectives are the categories of algebras for cellular monads/theories with respect to a class of maps determined by \mathcal{A} and \mathcal{B} .

2. Background on globular theories

2.1 Globular sets The *globe category* \mathbb{G} is freely generated by the graph

$$0 \begin{array}{c} \xrightarrow{\sigma_1} \\ \xrightarrow{\tau_1} \end{array} 1 \begin{array}{c} \xrightarrow{\sigma_2} \\ \xrightarrow{\tau_2} \end{array} \dots \begin{array}{c} \xrightarrow{\sigma_{n-1}} \\ \xrightarrow{\tau_{n-1}} \end{array} n-1 \begin{array}{c} \xrightarrow{\sigma_n} \\ \xrightarrow{\tau_n} \end{array} n \dots$$

subject to the relations $\sigma_n \circ \sigma_{n-1} = \sigma_n \circ \tau_{n-1}$ and $\tau_n \circ \sigma_{n-1} = \tau_n \circ \tau_{n-1}$. These relations ensure that there are precisely two maps $\sigma_{n,m} : \tau_{n,m} : n \rightrightarrows m$ for $m > n$. We abbreviate $\sigma_{n,m}$ and $\tau_{n,m}$ by σ and τ when the context is clear.

The presheaf category $[\mathbb{G}^{op}, \mathbf{Set}]$ is the category of *globular sets*. A globular set $A : \mathbb{G}^{op} \rightarrow \mathbf{Set}$ is specified by objects $A(n)$ together with morphisms

$$A(n) \begin{array}{c} \xrightarrow{s_n} \\ \xrightarrow{t_n} \end{array} A(n-1)$$

where we write $s_n = A(\sigma_n)$ and $t_n = A(\tau_n)$, or just s and t if the context is clear.


Elements of the set $A(n)$ are referred to as n -cells of A . A pair (x, y) of n -cells in A are said to be *parallel* if either $n = 0$ or if the equations $s_n x = s_n y$ and $t_n x = t_n y$ hold. A *lifting* for such a parallel pair (x, y) is an element $z \in A(n+1)$ such that $s_{n+1} z = x$ and $t_{n+1} z = y$. If each parallel pair of n -cells in A has a lifting (for each n) then the globular set A is said to be *contractible*.

The notions introduced in the preceding paragraph can be understood in terms of lifting properties. Firstly, an n -cell in A amounts to a morphism of globular sets $Yn \rightarrow A$ from the representable globular set $Yn = \mathbb{G}(-, n)$. Let $j_n : S(n) \hookrightarrow Y(n+1)$ be the globular subset of $Y(n+1)$ obtained by omitting the single $n+1$ -cell of $Y(n+1)$. This globular set has two distinct m -cells for all $m \leq n$ and none in higher dimensions. Now a parallel pair of n -cells (x, y) in A corresponds to a morphism $S(n) \rightarrow A$ and, moreover, the parallel pair admits a lifting just when the corresponding morphism $S(n) \rightarrow A$ admits an extension along $j_n : S(n) \hookrightarrow Y(n+1)$. Later on we will use the following well known expression of $S(n)$ as the coequaliser


$$Y(n-1) + Y(n-1) \begin{array}{c} \xrightarrow{\sigma^* + \sigma^*} \\ \xrightarrow{\tau^* + \tau^*} \end{array} Y(n) + Y(n) \xrightarrow{\langle j_n, j_n \rangle} S(n)$$

whose universal property captures the fact that $S(n)$ classifies parallel n -cells.


2.2 The category Θ_0 of globular cardinals The category of globular sets has a small dense subcategory Θ_0 , first described by Berger [5], whose objects have been termed *globular cardinals* by Street [21]. These include the representables—the n -globes Yn for each n —but also shapes such as the globular set with distinct cells as depicted below.



The globular cardinals can be parametrised in various ways, for instance using trees [4, 5]; following [19], we will use *tables of dimensions*—sequences $\vec{n} = (n_1, \dots, n_k)$ of natural numbers of odd length with $n_{2i-1} > n_{2i} < n_{2i+1}$. Given such a table \vec{n} and a functor $D : \mathbb{G} \rightarrow \mathcal{C}$, we obtain a diagram



whose colimit in \mathcal{C} , when it exists, will be written as $D(\vec{n})$, and called the *D-globular sum indexed by \vec{n}* . If all such colimits exist then we say that \mathcal{C} admits *D-globular sums* — let us emphasise that admitting globular sums is not a property of a category but of a category *under \mathbb{G}* . Dually a globular object $A : \mathbb{G}^{op} \rightarrow \mathcal{C}$ determines a diagram



whose limit, denoted $A(\vec{n})$, is called a *globular product*.

Consider $Y : \mathbb{G} \rightarrow [\mathbb{G}^{op}, \mathbf{Set}]$. Taking the full subcategory of $[\mathbb{G}^{op}, \mathbf{Set}]$ on the Y -globular sums yields the initial, up to equivalence, category under \mathbb{G} with globular sums. Θ_0 is a skeleton of this: we can view its objects as the tables of dimensions whilst $\Theta_0(\vec{n}, \vec{m}) = [\mathbb{G}^{op}, \mathbf{Set}](Y(\vec{n}), Y(\vec{m}))$. This gives a factorisation

$$\mathbb{G} \xrightarrow{D} \Theta_0 \xrightarrow{Y} [\mathbb{G}^{op}, \mathbf{Set}]$$

of the Yoneda embedding, in which $Dn = (n)$ on objects.

2.3 Globular theories and their models A *globular theory* $J : \Theta_0 \rightarrow \mathbb{T}$ is an identity on objects functor preserving globular sums. The *category* $\Theta_0\text{-Th}$ of *globular theories* is the full subcategory of Θ_0/\mathbf{Cat} containing the globular theories.

Given a globular theory $J : \Theta_0 \rightarrow \mathbb{T}$ its *category of models* $\text{Mod}(\mathbb{T}) \hookrightarrow [\mathbb{T}^{op}, \mathbf{Set}]$ is the full subcategory containing the globular product preserving functors. There is a forgetful functor $U^T : \text{Mod}(\mathbb{T}) \rightarrow [\mathbb{G}^{op}, \mathbf{Set}]$ given by restriction along $(J \circ D)^{op} : \mathbb{G}^{op} \rightarrow \mathbb{T}^{op}$. Furthermore a morphism $N : \mathbb{S} \rightarrow \mathbb{T}$ of globular theories induces, by restriction, a functor $N^* : \text{Mod}(\mathbb{T}) \rightarrow \text{Mod}(\mathbb{S})$ commuting with the forgetful functors to the category of globular sets.

The following proposition records a few basic results that we will need about globular theories and their models.

Proposition 2.1. *Let \mathbb{T} be a globular theory.*

1. *The forgetful functor $U^T : \text{Mod}(\mathbb{T}) \rightarrow [\mathbb{G}^{op}, \mathbf{Set}]$ is monadic and preserves filtered colimits. In particular $\text{Mod}(\mathbb{T})$ is locally finitely presentable.*
2. *Each representable $\mathbb{T}(-, \vec{n}) : \mathbb{T}^{op} \rightarrow \mathbf{Set}$ is a model of \mathbb{T} — namely the free T -model on the globular cardinal $Y\vec{n}$. Therefore we obtain a factorisation $K_T : \mathbb{T} \rightarrow \text{Mod}(\mathbb{T})$ of the Yoneda embedding Y_T through the full inclusion $I_T : \text{Mod}(\mathbb{T}) \hookrightarrow [\mathbb{T}^{op}, \mathbf{Set}]$.*
3. *Given a morphism $N : \mathbb{S} \rightarrow \mathbb{T}$ of globular theories the restriction functor $N^* : \text{Mod}(\mathbb{T}) \rightarrow \text{Mod}(\mathbb{S})$ has left adjoint $N_! = \text{Lan}_{K_S}(K_T \circ N) : \text{Mod}(\mathbb{S}) \rightarrow \text{Mod}(\mathbb{T})$.*

Proof. For monadicity in (1) see Proposition 6.3.6 of [2]. That U^T preserves filtered colimits follows easily from the fact that each globular cardinal $Y\vec{n}$ is finitely presentable as a globular set. It follows that the resulting monad $T = U^T F^T$ preserves filtered colimits. Since by [12, Satz 10.3] the category of algebras for a filtered colimit preserving monad on a locally finitely presentable category is again locally finitely presentable, this completes the proof of (1).

As representables send colimits to limits each representable is a \mathbb{T} -model, and the stated freeness property is simply an instance of the Yoneda lemma. This establishes (2). For (3) observe that since $\text{Mod}(\mathbb{T})$ is locally presentable, by (1), it is cocomplete. Therefore we can form the left Kan extension $L = \text{Lan}_{K_S}(K_T \circ N) : \text{Mod}(\mathbb{S}) \rightarrow \text{Mod}(\mathbb{T})$. It remains to prove that $L \dashv N^*$. Since the extension L is pointwise we have that $\text{Lan}_{K_S}(K_T \circ N) \cong \text{Lan}_{Y_S}(K_T \circ N) \circ \text{Mod}(\mathbb{S})(K_S -, 1)$, where the first component L' is the left Kan extension along the Yoneda embedding. Since the values of K_S are representable we have that $\text{Mod}(\mathbb{S})(K_S -, 1) \cong I_S$, where I_S is the full inclusion viewing models as presheaves. Hence we have natural isomorphisms

$$\text{Mod}(\mathbb{T})(LX, A) \cong \text{Mod}(\mathbb{T})(L' I_S X, A) \cong [\mathbb{S}^{op}, \mathbf{Set}](I_S X, \text{Mod}(\mathbb{T})(K_T N -, A))$$

with the second using the Kan adjunction $L' \dashv \text{Mod}(\mathbb{T})(K_T N -, 1)$. Since N^* acts by restriction we have $I_S \circ N^* \cong \text{Mod}(\mathbb{T})(K_T N -, 1)$ so that the right side above is naturally isomorphic to $[\mathbb{S}^{op}, \mathbf{Set}](I_S X, I_S N^* A)$. By fully faithfulness of I_S this in turn is naturally iso-

morphic to $\text{Mod}(\mathbb{S})(X, N^*A)$ and now the composite natural isomorphism $\text{Mod}(\mathbb{T})(LX, A) \cong \text{Mod}(\mathbb{S})(X, N^*A)$ proves the claim. \square

2.4 Contractible and cellular globular theories Let $J : \Theta_0 \rightarrow \mathbb{T}$ be a globular theory. A pair of morphisms

$$u, v : (n) \rightrightarrows \vec{m} \in \mathbb{T}$$

is said to be parallel if either $n = 0$ or $u \circ \sigma_n = v \circ \sigma_n$ and $u \circ \tau_n = v \circ \tau_n$. A *lifting* for the parallel pair consists of a morphism $h : (n + 1) \rightarrow \vec{m}$ such that $h \circ \sigma_{n+1} = u$ and $h \circ \rho_{n+1} = v$. If each parallel pair in \mathbb{T} has a lifting then \mathbb{T} is said to be *contractible*.

Remark 2.2. A parallel pair (u, v) in the globular theory \mathbb{T} as above is simply a parallel pair of n -cells in the globular set $\mathbb{T}(JD-, \vec{m}) : \mathbb{G}^{op} \rightarrow \mathbf{Set}$. In this way we see that \mathbb{T} is contractible just when the globular set $\mathbb{T}(JD-, \vec{m}) : \mathbb{G}^{op} \rightarrow \mathbf{Set}$ is contractible for each $\vec{m} \in \Theta_0$.

Consider a family \mathcal{I} of parallel pairs $u_i, v_i : (n_i) \rightrightarrows \vec{m}_i$ in the globular theory \mathbb{T} .¹ There exists a globular theory $\mathbb{T}_{\mathcal{I}}$ and morphism of globular theories

$$J_{\mathcal{I}} : \mathbb{T} \rightarrow \mathbb{T}_{\mathcal{I}}$$

obtained by *freely adjoining liftings* for the parallel pairs of \mathcal{I} . More precisely

1. For each $i \in I$ the parallel pair $u_i, v_i : (n_i) \rightrightarrows \vec{m}_i \in \mathbb{T}_{\mathcal{I}}$ is equipped with a lifting $\phi_i : (n_i + 1) \rightarrow \vec{m}_i$ and
2. Given a morphism of globular theories $Q : \mathbb{T} \rightarrow \mathbb{S}$ together with a lifting θ_i of the parallel pair (Qu_i, Qv_i) for each $i \in I$, there exists a unique morphism of globular theories $\bar{Q} : \mathbb{T}_{\mathcal{I}} \rightarrow \mathbb{S}$ such that $\bar{Q} \circ J_{\mathcal{I}} = Q$ and $\bar{Q}\phi_i = \theta_i$ for each $i \in I$.

These two properties characterise $\mathbb{T}_{\mathcal{I}}$ uniquely up to isomorphism under \mathbb{T} .

A *cellular* globular theory is, by definition, the colimit of a chain of globular theories

$$\Theta_0 = \mathbb{T}_0 \xrightarrow{J_1^0} \mathbb{T}_1 \longrightarrow \cdots \longrightarrow \mathbb{T}_n \xrightarrow{J_{n+1}^n} \mathbb{T}_{n+1} \xrightarrow{J_{\omega}^{n+1}} \mathbb{T}_{\omega} = \mathbb{T}$$

in which each $J_{n+1}^n : \mathbb{T}_n \rightarrow \mathbb{T}_{n+1}$ is of the form $J_{\mathcal{I}_{n+1}} : \mathbb{T}_n \rightarrow \mathbb{T}_{\mathcal{I}_{n+1}}$ for some family of parallel pairs \mathcal{I}_{n+1} of \mathbb{T}_n . We remark that cellularity in the above sense can be re-expressed as cellularity with respect to a set of maps — see Example 6.6.

A *coherator* is a cellular contractible globular theory and, by definition, a *Grothendieck weak ω -groupoid* is a model of a coherator. The homotopy theory of these structures has been studied in [19, 3] and more recently in [15, 17], though important questions remain open.

Now for \mathbb{T} cellular we have, on the level of models, a cochain

$$\begin{array}{ccccccc} \text{Mod}(\mathbb{T}) & & & & & & \\ & \searrow^{U_n^\omega} & & & \searrow^{U_0^\omega} & & \\ & & \text{Mod}(\mathbb{T}_n) & \longrightarrow & \cdots & \longrightarrow & \text{Mod}(\mathbb{T}_1) \xrightarrow{U_0^1} \text{Mod}(\Theta_0) \\ & \downarrow^{U_{n+1}^\omega} & & & & & \\ \cdots & \text{Mod}(\mathbb{T}_{n+1}) & \xrightarrow{U_{n+1}^n} & \text{Mod}(\mathbb{T}_n) & \longrightarrow & \cdots & \longrightarrow & \text{Mod}(\mathbb{T}_1) \xrightarrow{U_0^1} \text{Mod}(\Theta_0) \end{array}$$

where we write $U_n^m = (J_m^n)^*$ for all $n < m$. We refer to this as the *tower of models* of \mathbb{T} . Since taking categories of models sends colimits of globular theories to limits of categories over $\text{Mod}(\Theta_0)$ — see Section 5.3 of [11] — it follows that $\text{Mod}(\mathbb{T})$ is the limit of the above cochain of categories.

¹That is, a set I together with a parallel pair $u_i, v_i \in \mathbb{T}$ for each $i \in I$.

Faithfulness Conjecture 2.3 (Maltsiniotis). *Each of the connecting functors $J_m^n : \mathbb{T}_n \rightarrow \mathbb{T}_m$ defining a cellular globular theory \mathbb{T} is faithful.*

Remark 2.4. The conjecture is stated in Section 1.4 of Maltsiniotis [19] and in the thesis of Ara [2] as Conjecture 4.1.7. It is used in Proposition 6.7.15 and Corollary 6.7.16 of [2] to give an explicit description of the globular theory Θ_{BL} whose models coincide with the weak ω -categories of Batanin-Leinster [4, 18] — to be precise, the models of the initial globular operad with contraction.

Before moving on, let us identify the models of globular theories of the form $\mathbb{T}_{\mathcal{I}}$. To this end, we define a $(\mathbb{T}, \mathcal{I})$ -model (X, x) to consist of a \mathbb{T} -model X together with a morphism x_i for each $i \in I$ such that the diagram

$$\begin{array}{ccc}
 X(\vec{m}_i) & \xrightarrow{x_i} & X(n_i + 1) \\
 \searrow^{Xu_i} & & \swarrow_s \\
 & & X(n_i) \\
 \swarrow_{Xv_i} & & \nwarrow_t
 \end{array}$$

is serially commutative. The morphisms of $\text{Mod}(\mathbb{T}, \mathcal{I})$ commute with the given liftings. There is an evident forgetful functor $U : \text{Mod}(\mathbb{T}, \mathcal{I}) \rightarrow \text{Mod}(\mathbb{T})$.

Proposition 2.5. *There is an isomorphism of categories $K : \text{Mod}(\mathbb{T}, \mathcal{I}) \rightarrow \text{Mod}(\mathbb{T}_{\mathcal{I}})$ rendering commutative the diagram*

$$\begin{array}{ccc}
 \text{Mod}(\mathbb{T}_{\mathcal{I}}) & \xrightarrow{K} & \text{Mod}(\mathbb{T}, \mathcal{I}) \\
 \searrow_{J_{\mathcal{I}}^*} & & \swarrow_U \\
 & & \text{Mod}(\mathbb{T})
 \end{array}$$

Proof. The functor K sends a $\mathbb{T}_{\mathcal{I}}$ -model X to the $(\mathbb{T}, \mathcal{I})$ -model $(X \circ J_{\mathcal{I}}^{op}, X_{\phi})$ whose first component is $X \circ J_{\mathcal{I}}^{op} : \mathbb{T}^{op} \rightarrow \mathbb{T}_{\mathcal{I}}^{op} \rightarrow \mathbf{Set}$ and whose lifting function at $i \in I$ has value $X(\phi_i) : X(\vec{m}_i) \rightarrow X(n_i + 1)$. Clearly K commutes with the respective functors to $\text{Mod}(\mathbb{T})$. Hence it remains to show that it is bijective on objects and fully faithful. We establish these properties using endomorphism theories.

To this end, let $X \in \text{Mod}(\mathbb{T})$. By definition, the (relative) endomorphism theory of X is obtained as the (identity on objects/fully faithful)-factorisation of $X^{op} : \mathbb{T} \rightarrow \mathbf{Set}^{op}$ depicted on the left below

$$\begin{array}{ccc}
 \mathbb{T} & \xrightarrow{E_X} & \mathbb{E}nd(X) \\
 \searrow_{X^{op}} & & \downarrow_{M_X} \\
 & & \mathbf{Set}^{op}
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathbb{T} & \xrightarrow{E_X} & \mathbb{E}nd(X) & \xrightarrow{M_X} & \mathbf{Set}^{op} \\
 \downarrow F & \nearrow A & \searrow B^{op} & & \\
 \mathbb{S} & & & &
 \end{array}$$

Since X^{op} preserves globular sums so does E_X , so that $\mathbb{E}nd(X)$ becomes a globular theory on restricting E_X along $J : \Theta_0 \rightarrow \mathbb{T}$ and $E_X : \mathbb{T} \rightarrow \mathbb{E}nd(X)$ a morphism thereof. The functor M_X also preserves globular sums. The *universal property* of $\mathbb{E}nd(X)$ endowed with M_X is that, given a morphism of globular theories $F : \mathbb{T} \rightarrow \mathbb{S}$, composition with M_X induces a bijection between globular theory morphisms $A : \mathbb{S} \rightarrow \mathbb{E}nd(X)$ with $A \circ F = E_X$ and \mathbb{S} -model structures

$B : \mathbb{S}^{op} \rightarrow \mathbf{Set}$ such that $F^*B = X$. (This correspondence is indicated in the diagram above right and follows immediately from the fact that F , being identity on objects, and M_X , being fully faithful, are orthogonal.)

Now $\mathbb{E}nd(X)(\vec{n}, \vec{m}) = \mathbf{Set}(X\vec{m}, X\vec{n})$ with $E_X : \mathbb{T} \rightarrow \mathbb{E}nd(X)$ acting on morphisms as application of X . It follows that to equip X with the structure (X, x) of a $(\mathbb{T}, \mathcal{I})$ -model is equally to equip the morphism of theories $E_X : \mathbb{T} \rightarrow \mathbb{E}nd(X)$ with a lifting $(n_i + 1) \rightarrow \vec{m}_i$ of the parallel pair $(E_X u_i, E_X v_i)$ for each $i \in I$. Therefore by the universal property of $\mathbb{T}_{\mathcal{I}}$ there exists a unique morphism $\mathbb{T}_{\mathcal{I}} \rightarrow \mathbb{E}nd(X)$ extending E_X along $J_{\mathcal{I}}$; postcomposing with M_X gives the corresponding $\mathbb{T}_{\mathcal{I}}$ -model $K'(X, x)$. Spelling out the formulae involved we see that $KK'(X, x) = (X, x)$. To see that $K'K = 1$ we use the universal property of $\mathbb{T}_{\mathcal{I}}$.

The proof that K is bijective on arrows is almost identical, except that it uses endomorphism theories of *morphisms* of \mathbb{T} -models. A \mathbb{T} -model morphism $f : X \rightarrow Y$ corresponds to a globular product preserving functor $\mathbb{T} \rightarrow (\mathbf{Set}^2)^{op}$; now factoring it as (identity on objects/fully faithful) gives $E_f : \mathbb{T} \rightarrow \mathbb{E}nd(f)$, the (relative) endomorphism theory of $f : X \rightarrow Y$. We leave the details of this part to the reader. \square

3. Iterated algebraic injectives and models of cellular theories

3.1 Algebraic injectivity By a family $\mathcal{I} = (I, \alpha)$ of morphisms in \mathcal{C} we mean a set I and function $\alpha : I \rightarrow Arr(\mathcal{C})$.

Given such, an *algebraic \mathcal{I} -injective* is a pair (C, c) where $C \in \mathcal{C}$ together with extensions

$$\begin{array}{ccc} A_i & \xrightarrow{f} & C \\ \alpha_i \downarrow & \nearrow c(i,f) & \\ B_i & & \end{array}$$

for each lifting problem with $i \in I$. Morphisms $f : (C, c) \rightarrow (D, d)$ of algebraic injectives are morphisms $f : C \rightarrow D$ commuting with the given extensions. We write $\mathbb{I}nj(\mathcal{I})$ for the category of algebraic injectives and $V : \mathbb{I}nj(\mathcal{I}) \rightarrow \mathcal{C}$ for the forgetful functor to \mathcal{C} .

Notation 3.1. We sometimes write $|\mathcal{I}| = \{\alpha_i : i \in I\}$ for the underlying set of morphisms in \mathcal{C} . When we assert a property of morphisms of \mathcal{I} — for example that they are mono — this should be interpreted as a property of morphisms of $|\mathcal{I}|$.

Example 3.2. Let $\iota : 2 \hookrightarrow 3 \in \mathbf{Set}$ be the inclusion of the cardinal with 2 elements into the cardinal with 3. It is easy to see that $\mathbb{I}nj(\iota)$ is the category of magmas. More generally, consider a finitary signature $\Omega : \mathbb{N} \rightarrow \mathbf{Set}$ and the associated function $I_{\Omega} = \Sigma_{n \in \mathbb{N}} \Omega(n) \rightarrow Arr(\mathbf{Set})$ sending each element of $\Omega(n)$ to the inclusion $n \hookrightarrow n + 1$. Then $\mathbb{I}nj(I_{\Omega})$ is the category of Ω -algebras and, indeed, categories $\mathbb{I}nj(\mathcal{I})$ for families \mathcal{I} of morphisms amongst $\{n \hookrightarrow n + 1 : n \in \mathbb{N}\}$ are precisely the categories Ω -Alg of algebras for finitary signatures Ω .

Example 3.3. Consider the category of simplicial sets and $\mathcal{I} = \{\Lambda_n^k \rightarrow \Delta_n\}$ the set of horn inclusions. Then $\mathbb{I}nj(\mathcal{I})$ is the category of *algebraic Kan complexes* considered by Nikolaus [20].

The following is a standard result. The case dealing with a set, rather than a family, of morphisms is dealt with in the proof of Theorem 5 of [8] and the generalisation to a family of morphisms is trivial.

Proposition 3.4. *Let \mathcal{C} be a locally presentable category. Then $\mathbb{I}nj(\mathcal{I})$ is locally presentable and $V : \mathbb{I}nj(\mathcal{I}) \rightarrow \mathcal{C}$ an accessible strictly monadic right adjoint. Furthermore, if \mathcal{C} is locally finitely presentable and \mathcal{I} consists of morphisms with finitely presentable domains then $\mathbb{I}nj(\mathcal{I})$ is locally finitely presentable and V finitary.*

We now give the key example of algebraic injectivity pertaining to the study of cellular globular theories; namely, that $\mathbb{T}_{\mathcal{I}}$ -models are algebraic injectives in $Mod(\mathbb{T})$.

To see this, let \mathcal{I} be a family of parallel pairs in \mathbb{T} and consider the parallel pair $u_i, v_i : (n_i) \rightrightarrows \vec{m}_i$ for $i \in I$. This corresponds to the pair $u_{i*}, v_{i*} : \mathbb{T}(-, (n_i)) \rightrightarrows \mathbb{T}(-, \vec{m}_i)$ in $Mod(\mathbb{T})$, which — by Part 2 of Proposition 2.1 — we can equally write as a pair of maps

$$u_{i*}, v_{i*} : F(Y(n_i)) \rightrightarrows F(Y(\vec{m}_i))$$

between free \mathbb{T} -models on globular sets. That u_i and v_i are parallel in \mathbb{T} corresponds to the fact that we have a fork on the lower row below.

$$\begin{array}{ccc}
 F(Y(n_i - 1) + Y(n_i - 1)) & \xrightarrow[F(\tau^* + \tau^*)]{F(\sigma^* + \sigma^*)} & F(Yn_i + Yn_i) \xrightarrow{F\langle j_{n_i}, j_{n_i} \rangle} F(Sn_i) \\
 & & \searrow F\langle u_{i*}, v_{i*} \rangle \quad \downarrow (u_i, v_i) \\
 & & F(Y\vec{m}_i)
 \end{array}$$

Since the left adjoint F preserves coequalisers, the original pair u_i, v_i therefore corresponds to the factorisation

$$(u_i, v_i) : F(Sn_i) \rightarrow F(Y\vec{m}_i)$$

through the coequaliser. Now to give a \mathbb{T} -model X the structure of a $(\mathbb{T}, \mathcal{I})$ -model is to give for each $a : \mathbb{T}(-, \vec{m}_i) \rightarrow X$ and $i \in I$ an extension as below left

$$\begin{array}{ccc}
 FY(n_i) + FY(n_i) & \xrightarrow{\langle u_{i*}, v_{i*} \rangle} & FY(\vec{m}_i) & & F(Sn_i) & \xrightarrow{\langle u_i, v_i \rangle} & FY(\vec{m}_i) \\
 \langle \sigma_*, \tau_* \rangle \downarrow & & \downarrow a & & Fj_{n_i} \downarrow & & \downarrow a \\
 FY(n_i + 1) & \xrightarrow{?} & X & & FY(n_i + 1) & \xrightarrow{?} & X
 \end{array}$$

which, equivalently, is to give an extension rendering the rightmost square commutative. By the universal property of the pushout square

$$\begin{array}{ccc}
 F(Sn_i) & \xrightarrow{\langle u_i, v_i \rangle} & FY(\vec{m}_i) \\
 Fj_{n_i} \downarrow & & \downarrow k_i \\
 FY(n_i + 1) & \longrightarrow & FY(\vec{m}_i)_{u_i \simeq v_i}
 \end{array} \tag{1}$$

such are in bijective correspondence with extensions of a along k_i as below.

$$\begin{array}{ccc}
 FY(\vec{m}_i) & \xrightarrow{a} & X \\
 j_i \downarrow & \searrow & \uparrow \\
 FY(\vec{m}_i)_{u_i \simeq v_i} & &
 \end{array}$$

Accordingly we see that extensions of X to a $(\mathbb{T}, \mathcal{I})$ -model are in bijective correspondence with extensions of X to an algebraic \mathcal{I}^* -injective for the family \mathcal{I}^* of morphisms

$$I \rightarrow Arr(Mod(\mathbb{T})) : i \mapsto k_i : FY(n_i) \rightarrow FY(\vec{m}_i)_{u_i \simeq v_i} \tag{2}$$

Extending the above argument to morphisms in the evident manner, we obtain an isomorphism of categories $\text{Mod}(\mathbb{T}, \mathcal{I}) \cong \text{Inj}(\mathcal{I}^*)$ over $\text{Mod}(\mathbb{T})$. Composing this with the isomorphism of Proposition 2.5 we obtain one direction of:

Proposition 3.5. *The following coincide up to isomorphism over $\text{Mod}(\mathbb{T})$.*

- Forgetful functors $\text{Mod}(\mathbb{T}_{\mathcal{I}}) \rightarrow \text{Mod}(\mathbb{T})$ induced by morphisms of theories $\mathbb{T} \rightarrow \mathbb{T}_{\mathcal{I}}$ for \mathcal{I} a family of parallel pairs in \mathbb{T} ;
- Forgetful functors $\text{Inj}(\mathcal{I}) \rightarrow \text{Mod}(\mathbb{T})$ for \mathcal{I} a family of morphisms in $\text{Mod}(\mathbb{T})$, each of which is a pushout as in (1).

Proof. The reverse direction involves chasing backwards through the above constructions. \square

3.2 Iterated algebraic injectivity Let $I_1 \subseteq \text{Arr}(\mathcal{C})$ and $I_2 \subseteq \text{Arr}(\text{Inj}(\mathcal{I}_1))$ be a set of morphisms in the (non-algebraic) injectivity class $\text{Inj}(\mathcal{I}_1)$. Then $\text{Inj}(\mathcal{I}_2)$ is an injectivity class in \mathcal{C} itself — namely $\text{Inj}(\mathcal{I}_1 \cup \mathcal{I}_2)$ — and so iterating injectivity produces nothing new.

On the other hand *iterating algebraic injectivity* produces categories that cannot be obtained in a single step. In fact, the case of interest to us — Grothendieck weak ω -groupoids — uses ω iterations. We give the definition in its natural generality.

Definition 3.6. Let λ be an ordinal. By a tower of (algebraic) λ -injectives over \mathcal{C} we mean a cochain $\text{Inj}(\mathcal{I}) : (\lambda^+)^{op} \rightarrow \mathbf{CAT}$ such that

1. $\text{Inj}(\mathcal{I}_0) = \mathcal{C}$;
2. For each $n < \lambda$ the link map $V_n^{n+1} : \text{Inj}(\mathcal{I}_{n+1}) \rightarrow \text{Inj}(\mathcal{I}_n)$ is the forgetful functor from the category of algebraic injectives determined by a given family of maps \mathcal{I}_{n+1} of $\text{Inj}(\mathcal{I}_n)$.
3. the cochain is smooth: that is, at a limit ordinal $\gamma \leq \lambda$, we have $\text{Inj}(\mathcal{I}_\gamma) = \lim_{n < \gamma} \text{Inj}(\mathcal{I}_n)$.

The value of the cochain at $0 < \lambda \in \lambda^+$ gives the forgetful functor $V_0^\lambda : \text{Inj}(\mathcal{I}_\lambda) \rightarrow \mathcal{C}$ from the category of λ -injectives.

Thus 1-injectives are algebraic injectives in \mathcal{C} whilst 2-injectives are algebraic injectives in algebraic injectives, and so on.² By the general term iterated algebraic injectives we mean λ -injectives for some λ .

As mentioned, the primary structures of interest here are ω -injectives. A tower of ω -injectives is specified by a diagram

$$\begin{array}{ccccccc}
 & & \text{Inj}(\mathcal{I})_\omega & & & & \\
 & & \downarrow V_{n+1}^\omega & \searrow V_n^\omega & \searrow V_0^\omega & & \\
 \cdots & \text{Inj}(\mathcal{I}_{n+1}) & \xrightarrow{V_{n+1}^{n+1}} & \text{Inj}(\mathcal{I}_n) & \longrightarrow \cdots \longrightarrow & \text{Inj}(\mathcal{I}_1) & \xrightarrow{V_0^1} \mathcal{C}
 \end{array} \tag{3}$$

in which $\text{Inj}_\omega(\mathcal{I})$ is the limit of the sequence $\text{Inj}(\mathcal{I}_n)_{n < \omega}$.

Examples 3.7. All locally presentable categories are categories of iterated algebraic injectives in a power of \mathbf{Set} . This is established in the following sequence of examples.

1. If \mathcal{G} is a directed graph with set of objects O and directed edges E then the presheaf category $\mathbf{Set}^{\mathcal{G}}$ is a category of algebraic injectives $\text{Inj}(E)$ in \mathbf{Set}^O where $E \rightarrow \text{Arr}(\mathbf{Set}^O)$ sends $f : n \rightarrow m \in E$ to the coproduct inclusion $O(n, -) \rightarrow O(n, -) + O(m, -)$.

²We use the term λ -injective as opposed to algebraic λ -injective since there is no meaningful non-algebraic version of λ -injectivity beyond the case $\lambda = 1$.

2. If \mathcal{C} is a small category with underlying graph UC then it is easy to see that $[\mathcal{C}, \mathbf{Set}]$ is a small orthogonality class in $[UC, \mathbf{Set}]$. By Remark 4.4 of [1], each small orthogonality class in $[UC, \mathbf{Set}]$ is a small injectivity class — indeed, also a category of algebraic injectives in $[UC, \mathbf{Set}]$ (since the liftings are, in this case, forced to be unique.)
3. Since each locally presentable category is a small orthogonality class (thus a category of algebraic injectives) in a presheaf category, we conclude — on combining these examples — that each locally presentable category is a category of 3-injectives in a power of \mathbf{Set} .

Indeed the categories of iterated algebraic injectives in powers of \mathbf{Set} are precisely the locally presentable categories. If we only allow in each \mathcal{I}_n morphisms with finitely presentable domains we get precisely the locally finitely presentable categories. These claims follows from the following result.

Proposition 3.8. *Let $\mathbb{I}nj(\mathcal{I}) : (\lambda^+)^{op} \rightarrow \mathbf{CAT}$ be a tower of λ -injectives over \mathcal{C} . If \mathcal{C} is locally presentable then each category $\mathbb{I}nj(\mathcal{I}_n)$ in the tower is locally presentable and each connecting map $V_m^n : \mathbb{I}nj(\mathcal{I}_n) \rightarrow \mathbb{I}nj(\mathcal{I}_m)$ an accessible right adjoint. (If \mathcal{C} is locally finitely presentable and each \mathcal{I}_n consists of morphisms with finitely presentable domains then each $\mathbb{I}nj(\mathcal{I}_n)$ is locally finitely presentable and each connecting map a finitary right adjoint.)*

Proof. We prove the statement by transfinite induction on $n \leq \lambda$. The statement $n = 0$ is our assumption that \mathcal{C} is locally presentable. For $n = m + 1$ we have that $\mathbb{I}nj(\mathcal{I}_m)$ is locally presentable and V_k^m an accessible right adjoint for each $k < m$. By Proposition 3.4 then $\mathbb{I}nj(\mathcal{I}_{m+1})$ is locally presentable and V_m^{m+1} an accessible right adjoint — since each $V_k^{m+1} = V_k^m \circ V_m^{m+1}$ is then a composite of accessible right adjoints the claim holds. For n a limit ordinal the limit $\mathbb{I}nj(\mathcal{I}_n) = \mathbb{I}nj(\mathcal{I}_m)_{m < n}$ is, by Propositions A.1 and A.3, a bilimit. Since, by Theorem 2.18 of [6], the 2-category of locally presentable categories and accessible right adjoints is closed under bilimits in \mathbf{CAT} , the claim follows. The locally finitely presentable case follows similarly from the locally finitely presentable part of Proposition 3.4 and Theorem 2.17 of [6]. \square

We write $V_m^n : \mathbb{I}nj(\mathcal{I}_n) \rightarrow \mathbb{I}nj(\mathcal{I}_m)$ for the composite forgetful functor where $m < n \in \lambda^+$ and $F_m^n \dashv V_m^n$ for the adjoints. In the case $m = 0$ we write $V_n = V_0^n : \mathbb{I}nj(\mathcal{I}_n) \rightarrow \mathcal{C}$ for the composite and $F_n = F_0^n$ for its left adjoint.

The examples that we are interested in will arise from a more specialised context, in which we are provided with the data of a triple $(\mathcal{C}, \mathcal{A}, \mathcal{B})$ consisting of a locally presentable category \mathcal{C} , a set $\mathcal{A} \subseteq Ob(\mathcal{C})$ of objects and a family \mathcal{B} of morphisms of \mathcal{C} .

Now consider pushouts in $\mathbb{I}nj(\mathcal{I}_n)$ of the form

$$\begin{array}{ccc}
 F_n A & \xrightarrow{u} & F_n X \\
 F_n \alpha \downarrow & & \downarrow p \\
 F_n B & \xrightarrow{q} & P
 \end{array} \tag{4}$$

where $\alpha : A \rightarrow B \in |\mathcal{B}|$ and $X \in \mathcal{A}$.

Definition 3.9. The tower $\mathbb{I}nj(\mathcal{I}) : (\lambda^+)^{op} \rightarrow \mathbf{CAT}$ is a tower of $(\mathcal{A}, \mathcal{B})$ -iterated algebraic injectives if

- (†) for $n < \lambda$ each morphism of $|\mathcal{I}_{n+1}|$ is the pushout of a morphism $F_n \alpha : F_n A \rightarrow F_n B$ with $\alpha \in |\mathcal{B}|$ along a morphism $u : F_n A \rightarrow F_n X$ with $X \in \mathcal{A}$.

Proposition 3.10. *Let $\mathbb{I}nj(\mathcal{I}) : (\lambda^+)^{op} \rightarrow \mathbf{CAT}$ be a tower of $(\mathcal{A}, \mathcal{B})$ -iterated algebraic injectives. If each object of \mathcal{A} is finitely presentable and the family \mathcal{B} consists only of morphisms with finitely presentable domains, then each family \mathcal{I}_n has the same property.*

Proof. Inspecting the square (4) we see that each morphism in \mathcal{I}_n has source a free n -injective on a finitely presentable object. Therefore it suffices to show that each $F_n : \mathcal{C} \rightarrow \mathbb{I}nj(\mathcal{I}_n)$ preserves finitely presentable objects and, for this, it suffices to show that each right adjoint $V_n : \mathbb{I}nj(\mathcal{I}_n) \rightarrow \mathcal{C}$ preserves filtered colimits. This follows by transfinite induction, arguing as in Proposition 3.8. \square

Example 3.11. Let $\mathbb{F} \hookrightarrow \mathbf{Set}$ consist of the finite cardinals $0, 1, 2, \dots$ and $\mathcal{B} = \{! : 0 \rightarrow 1, ! : 2 \rightarrow 1\}$. From Example 3.2 $V : \mathbb{I}nj(\iota : 2 \rightarrow 3) \rightarrow \mathbf{Set}$ is the concrete category of magmas. Observe that we have a pushout square as below left.

$$\begin{array}{ccc}
 0 & \longrightarrow & 2 \\
 ! \downarrow & & \downarrow \iota \\
 1 & \longrightarrow & 3
 \end{array}
 \qquad
 \begin{array}{ccc}
 F2 & \xrightarrow{(t,s)} & F3 \\
 F! \downarrow & & \downarrow \\
 F1 & \longrightarrow & F3/\langle t = s \rangle
 \end{array}$$

Let F be the left adjoint of V . The terms $t = (xy)z, s = x(yz)$ in 3 variables are elements of $VF3$. Together, they correspond to a single map $\langle t, s \rangle : F2 \rightarrow F3$. By the universal property of the pushout $F3/\langle t = s \rangle$ each morphism $(a, b, c) : F3 \rightarrow A$ factors through $F3 \rightarrow F3/\langle t = s \rangle$ just when the equation $(ab)c = a(bc)$ holds for all $a, b, c \in A$. Thus we see that the category of associative magmas is a category of $(\mathbb{F}, \mathcal{B})$ -2-injectives.

Generalising this construction in the obvious way, each equational variety $(\Omega, E)\text{-Alg}$ is naturally a category of $(\mathbb{F}, \mathcal{B})$ -2-injectives. Conversely, each category of $(\mathbb{F}, \mathcal{B})$ -iterated algebraic injectives is an equational variety. One can prove this using sifted colimits and Beck’s theorem. A better argument, not using monadicity theorems, is provided by Theorem 6.3.

Recall the category Θ_0 of globular cardinals, and the set of morphisms $\mathcal{B} = \{S(n) \rightarrow Y(n+1) : n \in \mathbb{N}\}$ including the boundaries of representable globular sets. The main result of this section characterises models of cellular globular theories as (Θ_0, \mathcal{B}) - ω -injectives. We will state and prove this result in terms of cochains.

Both the towers of models of a cellular theory and of ω -injectives form ω^+ -cochains in \mathbf{CAT} where $\omega^+ = \{0 < 1, \dots < n < \dots < \omega\}$. These cochains share the properties of being smooth and *isofibrant*. Recall that a functor $W : \mathcal{A} \rightarrow \mathcal{B}$ is said to be an isofibration if given $A \in \mathcal{A}$ and an isomorphism $f : B \rightarrow WA \in \mathcal{B}$ there exists an isomorphism $f' : B' \rightarrow A$ with $Wf' = f$. Now a cochain X is said to be isofibrant if each $X_n \rightarrow X_m$ for $m < n$ is an isofibration of categories.

By an *equivalence* $F : X \rightarrow Y$ of cochains in \mathbf{CAT} we shall mean a natural transformation between cochains for which $F_n : X_n \rightarrow Y_n$ is an equivalence of categories for each n . If we restrict our attention to smooth isofibrant cochains then this yields the *correct* notion — in particular, it yields an *equivalence relation* on smooth isofibrant cochains (see Appendix A for a proof).

Theorem 3.12. *Up to equivalence of ω^+ -cochains, the towers of models of cellular globular theories and towers of (Θ_0, \mathcal{B}) - ω -injectives coincide.*

Proof. Consider the tower of models of a cellular globular theory \mathbb{T} . We will construct the equivalent tower of ω -injectives inductively. For the base case we use the canonical equivalence

$E_0 = U^{\Theta_0} : \text{Mod}(\mathbb{T}_0) = \text{Mod}(\Theta_0) \rightarrow [\mathbb{G}^{op}, \mathbf{Set}]$. Now suppose that we have a morphism of chains of length n as below

$$\begin{array}{ccccccc} \text{Mod}(\mathbb{T}_n) & \longrightarrow & \cdots & \longrightarrow & \text{Mod}(\mathbb{T}_1) & \longrightarrow & \text{Mod}(\mathbb{T}_0) \\ E_n \downarrow & & & & \downarrow E_1 & & \downarrow E_0 \\ \text{Inj}(\mathcal{I}_n, \alpha_n) & \longrightarrow & \cdots & \longrightarrow & \text{Inj}(\mathcal{I}_1, \alpha_1) & \longrightarrow & [\mathbb{G}^{op}, \mathbf{Set}] \end{array}$$

in which each E_j is an equivalence and each family \mathcal{I}_m satisfies (\dagger) for $m \leq n$. Since \mathbb{T} is cellular we have $\mathbb{T}_{n+1} = (\mathbb{T}_n)_{\mathcal{I}_{n+1}}$. Therefore by Proposition 3.5 we have an isomorphism $\text{Mod}(\mathbb{T}_{n+1}) \rightarrow \text{Inj}(\mathcal{I}_{n+1}, j_{n+1})$ making the left triangle below commute.

$$\begin{array}{ccccccc} \text{Mod}(\mathbb{T}_{n+1}) & \longrightarrow & \text{Inj}(\mathcal{I}_{n+1}, j_{n+1}) & \longrightarrow & \text{Inj}(\mathcal{I}_{n+1}, E_n j_{n+1}) & \longrightarrow & \text{Inj}(\mathcal{I}_{n+1}, \alpha_{n+1}) \\ & \searrow U_{n+1}^n & \downarrow & & \downarrow & \swarrow V_{n+1}^n & \\ & & \text{Mod}(\mathbb{T}_n) & \xrightarrow{E_n} & \text{Inj}(K_n) & & \end{array}$$

Here \mathcal{I}_{n+1} is of the form described in (2). Any equivalence of categories $W : \mathcal{A} \rightarrow \mathcal{B}$ lifts to an equivalence $\text{Inj}(S, \alpha) \rightarrow \text{Inj}(S, W\alpha)$ for (S, α) a family of morphisms in \mathcal{A} ; applying this to E_n and $(\mathcal{I}_{n+1}, j_{n+1})$ yields the equivalence in the central square above. The morphisms in the image of $E_n j_{n+1}$ do not satisfy (\dagger) on the nose — in particular, they have domains of the form $E_n F^{T_n} X$ for X a globular cardinal; however, composing these with the natural isomorphism $F_n X \cong E_n F^{T_n} X$ we obtain an isomorphic family $(\mathcal{I}_{n+1}, \alpha_{n+1})$ satisfying (\dagger) — this induces an isomorphism as in the triangle above right. We define E_{n+1} to be the composite equivalence on the top row above. Now since each E_n is an equivalence and since the cofiltered limits $\text{Mod}(\mathbb{T})$ and $\text{Inj}_\omega(K)$ are *bilimits* (see Propositions A.1 and A.3 of Appendix A) the induced map $E : \text{Mod}(\mathbb{T}) \rightarrow \text{Inj}_\omega(K)$ is an equivalence. This gives the desired equivalence of cochains. The reverse direction is similar. \square

The tower-free version is an immediate consequence.

Theorem 3.13. *Up to equivalence over $[\mathbb{G}^{op}, \mathbf{Set}]$, the categories of models of cellular globular theories and of (Θ_0, \mathcal{B}) - ω -injectives coincide.*

The above results mean that if we are only interested in the models of cellular globular theories, then we can equally work the semantic notion of (Θ_0, \mathcal{B}) -iterated algebraic injectives instead. The following result refines this to deal with cellular contractible theories (aka coherators) whose models are, by definition, the categories of Grothendieck weak ω -groupoids.

Theorem 3.14. *Up to equivalence over $[\mathbb{G}^{op}, \mathbf{Set}]$, the categories of models of coherators coincide with those categories of (Θ_0, \mathcal{B}) - ω -injectives having the property that each free ω -injective on a globular cardinal has a contractible underlying globular set.*

Proof. Consider the category of models of a cellular theory \mathbb{T} and the equivalence E with a category of (Θ_0, \mathcal{B}) - ω -injectives depicted below.

$$\begin{array}{ccc} \text{Mod}(\mathbb{T}) & \xrightarrow{E} & \text{Inj}_\omega(\mathcal{I}) \\ & \searrow U^T \quad \swarrow V_\omega & \\ & & [\mathbb{G}^{op}, \mathbf{Set}] \end{array}$$

Since E commutes with the forgetful functors it also commutes with the left adjoints, up to isomorphism. Thus the free ω -injective on each globular cardinal will be contractible just when the corresponding fact holds for the cellular globular theory \mathbb{T} : the free \mathbb{T} -model on each globular cardinal is contractible. So it remains to show that this last property holds just when \mathbb{T} is contractible. The free \mathbb{T} -model on a globular cardinal \vec{m} is simply $\mathbb{T}(-, \vec{m}) : \mathbb{T}^{op} \rightarrow \mathbf{Set}$ and this has underlying globular set the composite $\mathbb{T}(J \circ D-, \vec{m}) : \mathbb{G}^{op} \rightarrow \mathbf{Set}$. By Remark (2.2) each such globular set is contractible just when \mathbb{T} is so, as required. \square

Remark 3.15. One can tighten the correspondence of Theorem 3.12 by working with *concrete models* [11] of a globular theory \mathbb{T} . A concrete model of \mathbb{T} consists of a pair (X, A) where X is a globular set and $A : \mathbb{T}^{op} \rightarrow \mathbf{Set}$ an extension of $[\mathbb{G}^{op}, \mathbf{Set}](Y-, X) : \Theta_0^{op} \rightarrow \mathbf{Set}$ along J^{op} . The assignment $(X, A) \mapsto A$ forms the object part of an equivalence $\text{Mod}(\mathbb{T})_c \rightarrow \text{Mod}(\mathbb{T})$ between the categories of concrete and of ordinary models. Thus the two categories are essentially the same — however there is an isomorphism $\text{Mod}(\Theta_0)_c \cong [\mathbb{G}^{op}, \mathbf{Set}]$ rather than just an equivalence. A consequence is that the towers of concrete models of globular theories coincide, up to *isomorphism* of ω^+ -cochains, with the towers of (Θ_0, \mathcal{B}) - ω -injectives.

4. Free iterated algebraic injectives and cellularity

For \mathcal{C} a cofibrantly generated model category with generating trivial cofibrations \mathcal{I} one can define the category of *algebraically fibrant objects* over \mathcal{C} as $V : \mathbb{I}nj(\mathcal{I}) \rightarrow \mathcal{C}$. In Theorem 2.20 of [20] Nikolaus showed that if

- each trivial cofibration is a monomorphism and
- each morphism in \mathcal{I} has finitely presentable domain

then the model structure on \mathcal{C} can be right induced along V to a model structure on $\mathbb{I}nj(\mathcal{I})$ for which the adjunction $F \dashv V$ is a Quillen equivalence.³

The main point in establishing the model structure is to show that in pushouts of the form

$$\begin{array}{ccc} FA & \longrightarrow & X \\ F\alpha \downarrow & & \downarrow f \\ FB & \longrightarrow & Y \end{array}$$

with $\alpha \in \mathcal{I}$ the map $Vf : VX \rightarrow VY$ is a weak equivalence. That the resulting Quillen adjunction is a Quillen equivalence follows on showing that the unit

$$\eta_X : X \rightarrow VFX$$

is a weak equivalence for all X . Nikolaus achieved these two results by giving explicit, closely related, constructions of such pushouts and of free algebraically fibrant objects. In fact, Nikolaus did more than this — showing in fact that the sought for weak equivalences are *trivial cofibrations* — even \mathcal{I} -cellular maps.

This last point — a minor one at first appearance — obtains particular importance for us. The reason is that we will not be working with model structures and their weak equivalences, but simply a set of morphisms (or, if one prefers, the cofibrantly generated weak factorisation system they determine). In our setting we would like to obtain similar free constructions of iterated

³In fact, if \mathcal{C} is combinatorial, neither condition is required — see Theorem 15 of [8]. However for our present purposes we make use of these conditions and follow Nikolaus' original arguments closely.

algebraic injectives and pushouts. Therefore we begin by abstracting, in Section 4.1, the results of Nikolaus [20] to our setting. (These abstractions involve little originality since all of the core constructions described below in Section 4.1 are in *ibid.*) In Section 4.2 we extend these results to the iterated setting. This leads naturally to a proof of the faithfulness conjecture.

4.1 Free algebraic injectives and cellularity Let \mathcal{I} be a family of morphisms in a category \mathcal{C} . Let $Cell(\mathcal{I})$ denote the class of \mathcal{I} -cellular morphisms: a morphism is \mathcal{I} -cellular if it can be written as a transfinite composite of pushouts of small coproducts of maps in $|\mathcal{I}|$. It is well known and straightforward to show that $Cell(\mathcal{I})$ is stable under pushouts, coproducts and transfinite composition.

In what follows, we work in the context of cocomplete category \mathcal{C} equipped with a family of morphisms \mathcal{I} with finitely presentable domains such that $Cell(\mathcal{I}) \subseteq Mono$.

The constructions of algebraic injectives of interest to us presently are naturally seen as liftings of sinks along $V : \mathbb{I}nj(\mathcal{I}) \rightarrow \mathcal{C}$. Recall that given a set of objects $\{Y_j : j \in J\}$ a sink under Y consists of an object X and morphisms $\{f_j : Y_j \rightarrow X : j \in J\}$. Morphisms of the category $Sink(Y)$ of sinks commute with the coprojections from the Y_j in the evident manner. Given a functor $U : \mathcal{C} \rightarrow \mathcal{D}$ we obtain a functor

$$U_Y : Sink(Y) \rightarrow Sink(UY)$$

given by application of U . By definition, a U -semifinal lifting of a sink $S = (f_j : UY_j \rightarrow X : j \in J)$ is a representation of the functor $Sink(UY)(S, U_Y -) : Sink(Y) \rightarrow \mathbf{Set}$. Assuming, as is usually done, that U is faithful then this amounts to an object X_S of \mathcal{C} and morphism $\eta_S : X \rightarrow UX_S$ with the property that each $\eta_S \circ f_j : UY_j \rightarrow UX_S$ is in the image of U , and which is the initial such morphism.

We are interested in two constructions, which are special cases of semifinal liftings along $V : \mathbb{I}nj(\mathcal{I}) \rightarrow \mathcal{C}$ where $|J| = \emptyset$ or $|J| = 1$ respectively. These are

1. The free algebraic injective on $X \in \mathcal{C}$;
2. The semifinal lifting along V of a monomorphism $f : V(Y, y) \rightarrow X$.

The first case is the semifinal lifting of the empty sink over X whilst in the second case we view f as a 1-element sink. We treat the two cases together by constructing the semifinal lifting of a sink $S = \{f_j : V(Y_j, y_j) \rightarrow X : j \in J\}$ having $|J| \leq 1$ and each f_j a monomorphism. The semifinal lifting will be the colimit

$$\begin{array}{ccccccc}
 X_0 & \xrightarrow{p_1^0} & X_1 & \longrightarrow & \cdots & \longrightarrow & X_n & \xrightarrow{p_{n+1}^n} & X_{n+1} & \cdots \\
 & & & & & & & \searrow^{p_n} & \downarrow^{p_{n+1}} & \\
 & & & & & & & & & X_S
 \end{array}$$

of a chain in \mathcal{C} . We start by setting $X_0 = X$. For each n a subset

$$\mathcal{C}_n(A, X_n) \subseteq \mathcal{C}(A, X_n)$$

is specified and X_{n+1} is then defined as the pushout below

$$\begin{array}{ccc}
 \sum_{i \in I} \mathcal{C}_n(A_i, X_n).A_i & \longrightarrow & X_n \\
 \sum_{i \in I} \mathcal{C}_n(A_i, X_n).\alpha_i \downarrow & & \downarrow^{p_{n+1}^n} \\
 \sum_{i \in I} \mathcal{C}_n(A_i, X_n).B_i & \xrightarrow{x_{n+1}} & X_{n+1} .
 \end{array}$$

Using the coproduct inclusions, we see that X_{n+1} is the *universal object* equipped with a map $p_{n+1}^n : X_n \rightarrow X_{n+1}$ and liftings $x_{n+1}(r, i)$ for each pair $(i \in I, r \in \mathcal{C}_n(A_i, X_n))$, as depicted below.

$$\begin{array}{ccc} A_i & \xrightarrow{r} & X_n \\ \alpha_i \downarrow & & \downarrow p_{n+1}^n \\ B_i & \xrightarrow{x_{n+1}(r,i)} & X_{n+1} \end{array} \quad (5)$$

Observe also that each connecting map $p_{n+1}^n : X_n \rightarrow X_{n+1}$, is \mathcal{I} -cellular, and so a monomorphism.

Now let $\mathcal{C}_0(A, X_0) \subseteq \mathcal{C}(A, X_0)$ consist of those morphisms not factoring through a member f_j of the sink. For higher n we let $\mathcal{C}_n(A, X_n)$ be the set of arrows $A \rightarrow X_n$ that do not factor through $p_{n-1}^n : X_{n-1} \rightarrow X_n$.

We must equip X_S with the structure of an algebraic injective. To this end, consider $i \in I$ and $r : A_i \rightarrow X_S$. There are two cases.

If r factors through $p_0 \circ f_j : Y_j \rightarrow X \rightarrow X_S$ as r' , then this factorisation is unique since both p_0 and f_j are monic.

$$\begin{array}{ccccc} & & r & & \\ & & \curvearrowright & & \\ A_i & \xrightarrow{r'} & Y_j & \xrightarrow{p_0 \circ f_j} & X_S \\ \alpha_i \downarrow & & \nearrow y_j(r', i) & & \\ B_i & & & & \end{array}$$

We then define $x_S(r, i) = p_0 \circ f_j \circ y_j(r', i)$ as the composite filler depicted. Note that this definition is forced upon us by the requirement that $p_0 \circ f_j$ be a morphism of algebraic injectives.

Otherwise, since A_i is finitely presentable $r : A_i \rightarrow X_S$ factors through $p_n : X_n \rightarrow X_S$ for a smallest $n \in \mathbb{N}$. Again the factored morphism r' is unique since p_n is a monomorphism.

$$\begin{array}{ccccc} & & r & & \\ & & \curvearrowright & & \\ A_i & \xrightarrow{r'} & X_n & \xrightarrow{p_n} & X_S \\ \alpha_i \downarrow & & \downarrow p_{n+1}^n & & \nearrow p_{n+1} \\ B_i & \xrightarrow{x_{n+1}(r', i)} & X_{n+1} & & \end{array}$$

We define $x_S(r, i) = p_{n+1} \circ x_{n+1}(r', i)$ as the composite filler depicted.

Proposition 4.1. *Consider a cocomplete category \mathcal{C} equipped with a family of morphisms \mathcal{I} with finitely presentable domains such that $\text{Cell}(\mathcal{I}) \subseteq \text{Mono}$. Consider $V : \text{Inj}(\mathcal{I}) \rightarrow \mathcal{C}$.*

1. *The forgetful functor V has a left adjoint F and the unit $\eta_X : X \rightarrow VF X$ is \mathcal{I} -cellular for each $X \in \mathcal{C}$.*
2. *Given a monomorphism $f : V(Y, y) \rightarrow X$, its semifinal lifting $\eta_f : X \rightarrow V(X_f, x_f)$ exists and is \mathcal{I} -cellular.*

Proof. It remains to verify that the morphism $p_0 : X \rightarrow X_S = V(X_S, x_S)$ has the required universal property. To this end consider $k : X \rightarrow Z = V(Z, z)$ with each $k \circ f_j : Y_j \rightarrow Z$ a morphism of algebraic injectives. By the universal property of the colimit X_S , extensions of k as below left

$$\begin{array}{ccc} X_0 & & X_n \\ p_0 \downarrow & \searrow k & \downarrow p_{n+1}^n \\ X_S & \xrightarrow{k_S} & Z \\ & & \nearrow k_{n+1} \\ & & X_{n+1} \end{array}$$

are in bijection with families k_n satisfying the commutativity above right and having $k_0 = k$.

Using the case-by-case definition of the liftings x_S it is, moreover, straightforward to see that k_S is a morphism of algebraic injectives if and only if each $k \circ f_j : Y_j \rightarrow X \rightarrow X_S$ is so and when, furthermore, for all $i \in I$, $n \in \mathbb{N}$ and $r \in \mathcal{C}_n(A, X_n)$ the two composites below from B_i to Z coincide.

$$\begin{array}{ccc}
 A_i & \xrightarrow{r} & X_n & \xrightarrow{k_n} & Z \\
 \alpha_i \downarrow & & & \nearrow z(k_n \circ r, i) & \\
 B_i & & & &
 \end{array}
 \qquad
 \begin{array}{ccc}
 A_i & \xrightarrow{r} & X_n & \xrightarrow{k_n} & Z \\
 \alpha_i \downarrow & & p_{n+1}^n \downarrow & \nearrow k_{n+1} & \\
 B_i & \xrightarrow{x_{n+1}(r,i)} & X_{n+1} & &
 \end{array}$$

Now by the universal property of X_{n+1} as in (5) such a k_{n+1} is determined uniquely by k_n subject to the above equality — thus a family k_n satisfying the equality in (5) is uniquely determined by the component $k_0 = k$, and the claim follows. \square

Corollary 4.2. *Consider a cocomplete category \mathcal{C} equipped with a family of morphisms \mathcal{I} with finitely presentable domains such that $\text{Cell}(\mathcal{I}) \subseteq \text{Mono}$. Let $\alpha : A \rightarrow B$ be mono and consider $V : \text{Inj}(\mathcal{I}) \rightarrow \mathcal{C}$ with left adjoint F . The pushout*

$$\begin{array}{ccc}
 FA & \longrightarrow & X \\
 F\alpha \downarrow & & \downarrow f \\
 FB & \longrightarrow & Y
 \end{array}$$

exists and $Vf : VX \rightarrow VY$ is a composite of a pushout of α followed by an \mathcal{I} -cellular morphism.

Proof. Given the preceding result, this is a consequence of the following straightforward construction of pushouts in the context of semifinal liftings. Namely, consider an adjunction $F \dashv U : \mathcal{C} \rightarrow \mathcal{D}$ and a span as below left.

$$\begin{array}{ccc}
 FA & \xrightarrow{g} & X \\
 F\alpha \downarrow & & \\
 FB & &
 \end{array}
 \qquad
 \begin{array}{ccccc}
 A & \xrightarrow{\bar{g}} & UX & & \\
 \alpha \downarrow & & \downarrow p & \searrow Uk_p & \\
 B & \xrightarrow{q} & Y & \xrightarrow{\eta_p} & UY_p
 \end{array}
 \qquad
 \begin{array}{ccc}
 FA & \xrightarrow{g} & X \\
 F\alpha \downarrow & & \downarrow k_p \\
 FB & \xrightarrow{\eta_p \circ \bar{q}} & Y_p
 \end{array}$$

Let Y denote the denote the pushout of the corresponding span in \mathcal{D} , obtained by transposing through the adjunction, and Y_p the semi-final lifting of the pushout coprojection $p : UX \rightarrow Y$. Then Y_p is the pushout of the original span in \mathcal{C} , with coprojections as on the right above. \square

The following abstracts Theorem 2.20 of [20] away from model categories to the cellular setting.

Theorem 4.3. *Let \mathcal{C} be a cocomplete category and \mathcal{I} a family of morphisms in \mathcal{C} with finitely presentable domains such that $\text{Cell}(\mathcal{I}) \subseteq \text{Mono}$. Consider the forgetful functor $V : \text{Inj}(\mathcal{I}) \rightarrow \mathcal{C}$ with left adjoint F .*

1. Each unit component $\eta_X : X \rightarrow VFX$ is \mathcal{I} -cellular;
2. The forgetful functor $V : \text{Inj}(\mathcal{I}) \rightarrow \mathcal{C}$ sends $F\mathcal{I}$ -cellular morphisms to \mathcal{I} -cellular morphisms.
3. The left adjoint F is faithful.

It is a special case — taking $\mathcal{I} = \mathcal{B}$ — of the result that we require, which is stated and proven below. This is formulated with iteration in mind.

Theorem 4.4. *Let \mathcal{B} be a family of morphisms in the cocomplete category \mathcal{C} such that $\text{Cell}(\mathcal{B}) \subseteq \text{Mono}$. Let \mathcal{I} be a family of \mathcal{B} -cellular morphisms having finitely presentable domains, and consider the forgetful functor $V : \mathbb{I}nj(\mathcal{I}) \rightarrow \mathcal{C}$ with left adjoint F .*

1. *Each unit component $\eta_X : X \rightarrow VFX$ is \mathcal{B} -cellular;*
2. *The forgetful functor $V : \mathbb{I}nj(\mathcal{I}) \rightarrow \mathcal{C}$ sends $F\mathcal{B}$ -cellular morphisms to \mathcal{B} -cellular morphisms;*
3. *$\text{Cell}(F\mathcal{B}) \subseteq \text{Mono}$.*
4. *The left adjoint F is faithful.*

Proof. 1. By Proposition 4.1 $\eta_X : X \rightarrow VFX$ is \mathcal{I} -cellular. Since $\mathcal{I} \subseteq \text{Cell}(\mathcal{B})$, and $\text{Cell}(\mathcal{B})$ is closed under pushouts, coproducts and transfinite composition, we have that $\text{Cell}(\mathcal{I}) \subseteq \text{Cell}(\mathcal{B})$, whence η_X is also \mathcal{I} -cellular.

2. By Corollary 4.2 each pushout f of a morphism $F\alpha$ in $F\mathcal{B}$ is sent by V to a pushout of f (a \mathcal{B} -cellular morphism) followed by an \mathcal{I} -cellular, and hence \mathcal{B} -cellular, morphism. Since \mathcal{B} -cellular morphisms are closed under composition Vf is \mathcal{B} -cellular. Since the source and target of each $\alpha : A \rightarrow B$ in \mathcal{I} is finitely presentable V preserves filtered colimits, and so, transfinite composition. Thus $V(\text{Cell}(F\mathcal{B})) \subseteq \text{Cell}(\mathcal{B})$.

3. This follows from the preceding part on using that $\text{Cell}(\mathcal{B}) \subseteq \text{Mono}$ and that V , like any faithful functor, reflects monos.

4. From the first part, we have that each unit component $\eta_X : X \rightarrow VFX$ is \mathcal{B} -cellular and so monic. We have a commutative triangle

$$\begin{array}{ccc} \mathcal{C}(Y, X) & \xrightarrow{F_{Y,X}} & \mathbb{I}nj(\mathcal{I})(FY, FX) \\ & \searrow_{\eta_X \circ -} & \downarrow \cong \\ & & \mathcal{C}(Y, VFX) \end{array}$$

whose vertical morphism is the adjointness bijection. Since η_X is monic the diagonal function is injective, whence so is the horizontal $F_{Y,X}$, as required. \square

4.2 Free iterated algebraic injectives and cellularity The following is our iterated version of the preceding result. This is specialised to deal with $(\mathcal{A}, \mathcal{B})$ -iterated algebraic injectives in Theorem 4.7.

Theorem 4.5. *Let \mathcal{C} be locally presentable and \mathcal{B} a family of morphisms in \mathcal{C} such that $\text{Cell}(\mathcal{B}) \subseteq \text{Mono}$. Consider a tower $\mathbb{I}nj(\mathcal{I})$ of ω -injectives such that for each $n < \omega$ each morphism of \mathcal{I}_{n+1} is an $F_n\mathcal{B}$ -cellular morphism with finitely presentable domain. Consider $m < n \in \omega^+$.*

1. *The forgetful functor $V_m^n : \mathbb{I}nj(\mathcal{I}_n) \rightarrow \mathbb{I}nj(\mathcal{I}_m)$ sends $F_n\mathcal{B}$ -cellular morphisms to $F_m\mathcal{B}$ -cellular morphisms.*
2. *The unit component $X \rightarrow V_m^n F_m^n X$ is $F_m\mathcal{B}$ -cellular for each $X \in \mathbb{I}nj(\mathcal{I}_m)$.*
3. *$\text{Cell}(F_n\mathcal{B}) \subseteq \text{Mono}$.*
4. *The left adjoint F_m^n is faithful.*

Proof. For finite n the stated properties hold by inductive application of Theorem 4.4 since, by that result, if $\text{Cell}(F_n\mathcal{B}) \subseteq \text{Mono}$ then $\text{Cell}(F_{n+1}\mathcal{B}) = \text{Cell}(F_n^{n+1}F_n\mathcal{B}) \subseteq \text{Mono}$ too. It remains to consider the case $n = \omega$ and $m < n$. In fact the case of general m follows from the case $m = 0$ since $\mathbb{I}nj_\omega(\mathcal{I})$ is equally the limit of the cochain $\mathbb{I}nj(\mathcal{I}_k)$ having $k \geq m$.

Therefore it remains to prove the three claims when $n = \omega$ and $m = 0$. Just as in the proof of Theorem 4.4, these will follow on showing that given a sink $S = \{f_j : V_0^\omega Y_j \rightarrow X : j \in J\}$ with

at most one member, a mono, its semifinal lifting exists and has unit $X \rightarrow V_0^\omega X_\omega$ a \mathcal{B} -cellular morphism. We begin by forming the semifinal lifting of the sink along V_0^1 , as depicted on the left below.

$$\begin{array}{ccc}
 V_0^\omega Y_j \xrightarrow{f_j=f_{j,0}} X & & V_n^\omega Y_j \xrightarrow{f_{j,n}} X_n \\
 \searrow V_0^1 f_{j,1} & \downarrow \eta_1^0 & \searrow V_n^{n+1} f_{j,n+1} & \downarrow \eta_{n+1}^n \\
 & V_0^1 X_1 & & V_n^{n+1} X_{n+1}
 \end{array}$$

By Theorem 4.3 the unit η_1^0 is \mathcal{B} -cellular. We now iterate this, forming the V_n^{n+1} -semifinal lifting of $f_{j,n}$ as on the right above. Note that this diagram lives in $\text{Inj}(\mathcal{I}_n)$. Arguing inductively, using the established case $n = m + 1$, we see that each $f_{j,n}$ is mono and that each η_{n+1}^n is $F_n \mathcal{B}$ -cellular. Now form the colimit of the following chain $(D_n)_{n \in \mathbb{N}}$ in \mathcal{C} .

$$\begin{array}{ccccccc}
 X & \xrightarrow{\eta_1^0} & V_0^1 X_1 & \longrightarrow & \cdots & \longrightarrow & V_0^n X_n & \xrightarrow{V_0^n \eta_{n+1}^n} & V_0^{n+1} X_{n+1} & \cdots \\
 & & & & & & & \searrow \eta_n & & \swarrow \eta_{n+1} \\
 & & & & & & & & \text{col}(D) &
 \end{array}$$

Since η_{n+1}^n is $F_n \mathcal{B}$ -cellular its underlying map $V_0^{n+1} \eta_{n+1}^n$ is \mathcal{B} -cellular. Thus the transfinite composite $\eta_0 : X \rightarrow \text{col}(D)$ is \mathcal{B} -cellular.

Observe that $\text{col}(D)$ is equally the colimit of the restricted diagram $(D_k)_{k \geq n}$, which lies in the image of V_0^n . Since V_0^n creates filtered colimits it follows that $\text{col}(D)$ obtains the structure of an n -injective $\text{col}(D)_n$, unique such that each cocone projection $\eta_k : V_n^k X_k \rightarrow \text{col}(D)_n$ is a morphism of $\text{Inj}(\mathcal{I}_n)$ for $m > n$. Since each V_n^{n+1} creates filtered colimits $V_n^{n+1} \text{col}(D)_{n+1} = \text{col}(D)_n$, so that the sequence $X_\omega = (\text{col}(D)_n)_{n \in \mathbb{N}}$ is an object of $\text{Inj}(\mathcal{I}_\omega)$.

Now consider the composite $\eta_0 : X \rightarrow \text{col}(D) = V_0^\omega X_\omega$. Then $\eta_0 \circ f_j = \eta_n \circ f_{j,n}$, both of whose components are morphisms of n -injectives. Thus $\eta_0 \circ f_j$ is a morphism of n -injectives for each n , and so a morphism of ω -injectives. For the universal property, consider $g : X \rightarrow V_0^\omega Z$ having $g \circ f_j$ a morphism of $\text{Inj}(\mathcal{I}_\omega)$.

$$\begin{array}{ccccc}
 V_0^\omega Y_j & \xrightarrow{f_j} & X & \xrightarrow{\eta_n^0} & V_0^n X_n & \xrightarrow{\eta_n} & V_0^\omega X_\omega \\
 & & \searrow g & & \downarrow g_n & & \swarrow g_\omega \\
 & & & & V_0^\omega Z & &
 \end{array}$$

Successively applying the universal properties of the semifinal liftings we obtain a unique morphism g_n of n -injectives satisfying $g_n \eta_n^0 = g_0$. These induce a unique map $g_\omega : V_0^\omega X_\omega \rightarrow V_0^\omega Z$ commuting with the g_n . Now to say that g_ω is a morphism of ω -injectives is equivalent to asking that its restriction along each η_n , the g_n , is a morphism of n -injectives; thus the unique extension g_ω is a morphism of ω -injectives, as required. \square

Remark 4.6. As a special case of the above we obtain the construction of free ω -injectives as colimits of chains of the form

$$X \longrightarrow V_0^1 F_0^1 X \longrightarrow \cdots \longrightarrow V_0^n F_0^n X \xrightarrow{V_0^n \eta F_0^n X} V_0^{n+1} F_0^{n+1} X \longrightarrow X_\omega .$$

Let us also state the version for $(\mathcal{A}, \mathcal{B})$ - ω -injectives. This follows immediately from Theorem 4.5 and Proposition 3.10.

Theorem 4.7. *Let \mathcal{C} be locally presentable, \mathcal{A} a set of finitely presentable objects in \mathcal{C} and \mathcal{B} a family of morphisms with finitely presentable domains in \mathcal{C} such that $\text{Cell}(\mathcal{B}) \subseteq \text{Mono}$. Given a tower $\mathbb{I}\text{nj}(\mathcal{I}_n)_{n \in \omega^+}$ of $(\mathcal{A}, \mathcal{B})$ - ω -injectives consider $n, m \in \omega^+$ with $m < n$.*

1. *The forgetful functor $V_m^n : \mathbb{I}\text{nj}(\mathcal{I}_n) \rightarrow \mathbb{I}\text{nj}(\mathcal{I}_m)$ sends $F_n \mathcal{B}$ -cellular morphisms to $F_m \mathcal{B}$ -cellular morphisms.*
2. *The unit component $X \rightarrow V_m^n F_m^n X$ is $F_m \mathcal{B}$ -cellular for each $X \in \mathbb{I}\text{nj}(\mathcal{I}_m)$.*
3. *For each n we have $\text{Cell}(F_n \mathcal{B}) \subseteq \text{Mono}$.*
4. *The left adjoint F_m^n is faithful.*

5. Cellular globular theories and the faithfulness conjecture

Consider a cellular globular theory

$$\Theta_0 = \mathbb{T}_0 \xrightarrow{J_1^0} \mathbb{T}_1 \longrightarrow \cdots \longrightarrow \mathbb{T}_n \xrightarrow{J_{n+1}^n} \mathbb{T}_{n+1} \xrightarrow{J_{\omega}^{n+1}} \mathbb{T}_{\omega} = \mathbb{T}$$

and its associated tower of models

$$\cdots \text{Mod}(\mathbb{T}_{n+1}) \xrightarrow{U_n^{n+1}} \text{Mod}(\mathbb{T}_n) \longrightarrow \cdots \longrightarrow \text{Mod}(\mathbb{T}_1) \xrightarrow{U_0^1} \text{Mod}(\Theta_0).$$

Composing with the equivalence $U^{\Theta_0} : \text{Mod}(\Theta_0) \rightarrow [\mathbb{G}^{op}, \mathbf{Set}]$ we obtain for $m < n \leq \omega$ a commutative triangle

$$\begin{array}{ccc} \text{Mod}(\mathbb{T}_n) & \xrightarrow{U_m^n} & \text{Mod}(\mathbb{T}_m) \\ & \searrow U^{T_n} & \downarrow U^{T_m} \\ & & [\mathbb{G}^{op}, \mathbf{Set}] \end{array} .$$

We write $F_m^n = (J_n^m)! : \text{Mod}(\mathbb{T}_m) \rightarrow \text{Mod}(\mathbb{T}_n)$ for the left adjoint to the forgetful functor and also $F_0^n = F^n$.

Theorem 5.1. *Given a cellular globular theory \mathbb{T} as above*

1. *The forgetful functor $U_m^n : \text{Mod}(\mathbb{T}_n) \rightarrow \text{Mod}(\mathbb{T}_m)$ sends $F_n \mathcal{B}$ -cellular morphisms to $F_m \mathcal{B}$ -cellular morphisms.*
2. *The unit component $X \rightarrow U_m^n F_m^n X$ is $F_m \mathcal{B}$ -cellular for each $X \in \text{Mod}(\mathbb{T}_m)$.*
3. *$\text{Cell}(F_n \mathcal{B}) \subseteq \text{Mono}$.*
4. *The left adjoint F_m^n is faithful.*

Proof. By Theorem 3.12 there is a (Θ_0, \mathcal{B}) -tower of ω -injectives and cochain equivalence as depicted below.

$$\begin{array}{ccccc} \text{Mod}(\mathbb{T}_n) & \xrightarrow{U_m^n} & \text{Mod}(\mathbb{T}_m) & \xrightarrow{U_0^m} & \text{Mod}(\Theta_0) \\ \downarrow E_n & & \downarrow E_m & \searrow U^{T_m} & \downarrow E_0 \\ \mathbb{I}\text{nj}(\mathcal{I}_n) & \xrightarrow{V_{n,m}} & \mathbb{I}\text{nj}(\mathcal{I}_m) & \xrightarrow{V_m} & [\mathbb{G}^{op}, \mathbf{Set}] \end{array}$$

By virtue of the cochain equivalence, we must show that the corresponding facts to (1),(2) and (3) hold for the tower of ω -injectives. This is precisely the content of Theorem 4.7 on observing that its hypotheses — each object of Θ_0 is finitely presentable as a globular set, each morphism of \mathcal{B} has finitely presentable domain and codomain and $\text{Cell}(\mathcal{B}) = \text{Mono}$ — are satisfied. \square

Theorem 5.2 (The faithfulness conjecture). *Each connecting map $J_n^m : \mathbb{T}_m \rightarrow \mathbb{T}_n$ defining a cellular globular theory \mathbb{T} is faithful.*

Proof. By Proposition 2.1 we have $F_m^n = \text{Lan}_{K_{T_m}}(K_{T_n} \circ J_n^m) : \text{Mod}(\mathbb{T}_m) \rightarrow \text{Mod}(\mathbb{T}_n)$ where K_{T_m} and K_{T_n} are the restricted Yoneda embeddings for \mathbb{T}_m and \mathbb{T}_n respectively. Since K_{T_m} is fully faithful it follows that we have a natural isomorphism in the square below.

$$\begin{array}{ccc}
 \mathbb{T}_m & \xrightarrow{J_n^m} & \mathbb{T}_n \\
 K_{T_m} \downarrow & \cong & \downarrow K_{T_n} \\
 \text{Mod}(\mathbb{T}_m) & \xrightarrow{F_m^n} & \text{Mod}(\mathbb{T}_n)
 \end{array}$$

Now both vertical morphisms are fully faithful whilst F_m^n is faithful by Theorem 5.1. Therefore J_n^m is faithful too. □

6. Cellular monads and cellular theories

In the present section, we briefly describe a generalisation of Theorem 3.12, which captures the models of cellular globular theories as ω -injectives. We will present our results in the general setting of [11]. In particular, our base \mathcal{E} will be *enriched* in a symmetric monoidal locally presentable category \mathcal{V} .

Given a family \mathcal{I} of morphisms in \mathcal{E} , we can form the *enriched category of algebraic injectives*, an object of which consists of $E \in \mathcal{E}$ together with a section $e(i, -) : \mathcal{E}(X_i, E) \rightarrow \mathcal{E}(Y_i, E)$ of $\mathcal{E}(\alpha_i, E) : \mathcal{E}(Y_i, E) \rightarrow \mathcal{E}(X_i, E)$ for each $i \in I$. As an enriched category $\mathbb{I}nj(\mathcal{I})$ can be encoded as the pullback:

$$\begin{array}{ccc}
 \mathbb{I}nj(\mathcal{I}) & \longrightarrow & \text{SE}[\mathcal{I}, \mathcal{V}] \\
 V \downarrow & & \downarrow \\
 \mathcal{E} & \xrightarrow{K} & \text{Arr}[\mathcal{I}, \mathcal{V}] .
 \end{array}$$

Here the right leg $\text{SE}[\mathcal{I}, \mathcal{V}] \rightarrow \text{Arr}[\mathcal{I}, \mathcal{V}]$ is the forgetful functor from the enriched category of *split epimorphisms* in $[\mathcal{I}, \mathcal{V}]$ to the enriched category of arrows therein, and K the \mathcal{V} -functor sending C to the family $(\mathcal{E}(\alpha_i, E) : \mathcal{E}(Y_i, E) \rightarrow \mathcal{E}(X_i, E))_{i \in I}$.

Example 6.1. Consider $\mathcal{V} = \mathbf{Cat}$ and $\mathcal{E} = \mathbf{Cat}$ as a 2-category. Building on Example 3.2, let $\iota : 2 \rightarrow 3$ be the inclusion of the discrete category on two objects to the discrete category on three. An object of $\mathbb{I}nj(\iota)$ is a category C equipped with a functor $C^2 \rightarrow C$ — in particular, $\mathbb{I}nj(\iota)$ is the 2-category of magmas in \mathbf{Cat} .

Now \mathcal{E} itself will be a locally presentable \mathcal{V} -category, which comes equipped with a small dense full sub \mathcal{V} -category $\mathcal{A} \hookrightarrow \mathcal{E}$. This is the basic setting of [11]. It is not hard to see — for instance, using Lemma 4 of *ibid.* — that $\mathbb{I}nj(\mathcal{I})$ is itself a locally presentable enriched category, whose enriched forgetful functor V to \mathcal{E} is itself a right adjoint.

In this setting one also has a good notion of \mathcal{A} -theory — capturing the globular theories when $\mathcal{A} = \Theta_0$ — and of \mathcal{A} -nervous monad. The two notions are equivalent by Theorem 17 of *ibid.* In the present section we prove our results using monads because their relationship with signatures, of central importance here, is a little cleaner to express. We also consider these results from the perspective of \mathcal{A} -theories at the end of this section.

Following *ibid.* the category $\mathbf{Sig}_{\mathcal{A}}(\mathcal{E})$ of *signatures* is the *ordinary* category $[ob\mathcal{A}, \mathcal{E}]$. Given $\Omega \in \mathbf{Sig}_{\mathcal{A}}(\mathcal{E})$ an Ω -algebra consists of an object $E \in \mathcal{E}$ together with a morphism $\mathcal{E}(A, E) \rightarrow \mathcal{E}(\Omega A, E)$ for each $A \in \mathcal{A}$. Ω -algebras, together with the natural structure on homs, form an enriched category $Alg(\Omega)$. Now there is a forgetful functor $U : \mathbf{Mnd}(\mathcal{E}) \rightarrow \mathbf{Sig}_{\mathcal{A}}(\mathcal{E})$ sending an enriched monad T to the family $(TA)_{A \in \mathcal{A}}$. By Theorem 36 of [11] this has a left adjoint F . Furthermore by the proof of Proposition 53 of *ibid.*, the \mathcal{V} -category of $F\Omega$ -algebras is, up to isomorphism over \mathcal{E} , the \mathcal{V} -category $Alg(\Omega)$ just described.

Now although the category of monads is not typically cocomplete it does admit all colimits of free monads on signatures. In fact the colimit closure $\mathbf{Mnd}_{\mathcal{A}}(\mathcal{E})$ of such free monads is extremely well behaved: it is locally presentable, monadic over $\mathbf{Sig}_{\mathcal{A}}(\mathcal{E})$ and contains exactly the so-called *\mathcal{A} -nervous monads*. For more on nervous monads and proofs of the above claims see [11].

The key signatures for us are the following ones. Namely, given a pair $(A \in \mathcal{A}, X \in \mathcal{E})$ we define the signature $A|X : ob\mathcal{A} \rightarrow \mathcal{E}$ to have value X at A and the initial object \emptyset otherwise. Thus a $A|X$ -algebra consists of an object E together with a single morphism $\mathcal{E}(A, E) \rightarrow \mathcal{E}(X, E)$. At $f : X \rightarrow Y$ we have the map $A|f : A|X \rightarrow A|Y$ with value f at A .

Evidently there is a natural isomorphism

$$\mathbf{Sig}_{\mathcal{A}}(\mathcal{E})(A|X, \Omega) \cong \mathcal{E}(X, \Omega(A)) \quad .$$

Given the morphism of signatures $A|f : A|X \rightarrow A|Y$ we can form the corresponding morphism $F(A|f)$ of monads; now a pushout of a coproduct of such maps

$$\begin{array}{ccc} F(\sum_{i \in I} A_i | X_i) & \xrightarrow{\langle t_i \rangle_{i \in I}} & T \\ F(\sum_{i \in I} A_i | f_i) \downarrow & & \downarrow P_{\mathcal{I}} \\ F(\sum_{i \in I} A_i | Y_i) & \longrightarrow & T_{\mathcal{I}} \end{array}$$

is specified by a family of triples $(f_i : X_i \rightarrow Y_i, t_i : F(A_i | X_i) \rightarrow T, A_i)_{i \in I}$. What are the algebras for the pushout $T_{\mathcal{I}}$? Since, by Proposition 28 of *ibid.*, semantics sends colimits to limits we have a pullback square as on the left below.

$$\begin{array}{ccccc} Alg(T_{\mathcal{I}}) & \longrightarrow & Alg(F(\sum_{i \in I} A_i | Y_i)) & \longrightarrow & Alg(\sum_{i \in I} A_i | Y_i) \\ Alg(P_{\mathcal{I}}) \downarrow & & \downarrow & & \downarrow Alg(\sum_{i \in I} A_i | f_i) \\ Alg(T) & \longrightarrow & Alg(F(\sum_{i \in I} A_i | X_i)) & \longrightarrow & Alg(\sum_{i \in I} A_i | X_i) \end{array}$$

Since we have horizontal isomorphisms in the right square above, the outer square is a pullback. Now under the isomorphisms

$$\mathbf{Mnd}(\mathcal{E})(F(A_i | X_i), T) \cong \mathbf{Sig}_{\mathcal{A}}(\mathcal{E})(A_i | X_i, UT) \cong \mathcal{E}(X_i, TA_i)$$

the map $t : F(A_i | X_i) \rightarrow T$ bijectively corresponds to a morphism $\bar{t} : X_i \rightarrow TA_i$. Furthermore a $\sum_{i \in I} A_i | X_i$ -algebra on $E \in \mathcal{E}$ is specified by an $A_i | X_i$ -algebra structure on E for each $i \in I$. In these terms, the horizontal composite functor on the bottom row of the above diagram sends the T -algebra (E, e) to the $\sum_{i \in I} A_i | X_i$ -algebra whose value at $i \in I$ is the morphism

$$\mathcal{E}(A_i, E) \xrightarrow{T} \mathcal{E}(TA_i, TE) \xrightarrow{\mathcal{E}(\bar{t}_i, e)} \mathcal{E}(X_i, E) \quad .$$

Therefore a $T_{\mathcal{I}}$ -algebra amounts to a T -algebra (E, e) , together with a lifting

$$\begin{array}{ccc} \mathcal{E}(A_i, E) & \xrightarrow{?} & \mathcal{E}(Y_i, E) \\ T \downarrow & & \downarrow \mathcal{E}(f_i, E) \\ \mathcal{E}(TA_i, TE) & \xrightarrow{\mathcal{E}(\bar{t}_i, e)} & \mathcal{E}(X_i, E) \end{array} \tag{6}$$

for each $i \in I$. Morphisms of $T_{\mathcal{I}}$ -algebras amount to morphisms of T -algebras commuting with such liftings in the evident sense, and they form the morphisms of a \mathcal{V} -enriched category in a natural way.

Now transposing \bar{t} through the isomorphism $\mathcal{E}(X_i, TA_i) \cong \text{Alg}(T)(F^T X_i, F^T A_i)$ to a map t_i^* we may form the following pushout

$$\begin{array}{ccc} F^T X_i & \xrightarrow{t_i^*} & F^T A_i \\ F^T f_i \downarrow & & \downarrow p_i \\ F^T Y_i & \longrightarrow & P_i \end{array} \tag{7}$$

in the enriched category of T -algebras. We thereby obtain a corresponding family $(p_i)_{i \in I}$ of morphisms in $\text{Alg}(T)$. Using the universal property of the pushout we see that to equip (E, e) with the structure of an algebraic \mathcal{I} -injective is equally to give a lifting as on the left below

$$\begin{array}{ccc} \text{Alg}(T)(F^T A_i, (E, e)) & \xrightarrow{?} & \text{Alg}(T)(F^T Y_i, (E, e)) & \mathcal{E}(A_i, E) & \xrightarrow{?} & \mathcal{E}(Y_i, E) \\ & \searrow \text{Alg}(T)(t_i^*, (E, e)) & \downarrow \text{Alg}(T)(F^T f_i, 1) & & \searrow \mathcal{E}(\bar{t}_i, e) \circ T & \downarrow \mathcal{E}(f_i, 1) \\ & & \text{Alg}(T)(F^T X_i, (E, e)) & & & \mathcal{E}(X_i, E) \end{array}$$

for each member of \mathcal{I} . Transposing through the adjunction $F^T \dashv U^T$ and using that the transpose of t_i^* is $\mathcal{E}(\bar{t}_i, e) \circ T$ as in (6) we conclude that this is precisely to equip (E, e) with the structure of a $T_{\mathcal{I}}$ -algebra. With a similar extension to hom-objects we obtain one direction of:

Proposition 6.2. *The following coincide up to isomorphism over $\text{Alg}(T)$.*

- Forgetful functors $\text{Alg}(T_{\mathcal{I}}) \rightarrow \text{Alg}(T)$ induced by morphisms of $T \rightarrow T_{\mathcal{I}}$ where $T \in \mathbf{Mnd}_{\mathcal{A}}(\mathcal{E})$.
- Forgetful functors $\mathbb{I}nj(\mathcal{I}) \rightarrow \text{Alg}(T)$ for \mathcal{I} a family of morphisms in $\text{Alg}(T)$, each of which is a pushout as in (7).

Proof. The reverse direction involves chasing backwards through the above constructions. □

Given a family \mathcal{B} of morphisms of \mathcal{E} let

$$F(\mathcal{A}|\mathcal{B}) = (F(A|f_i) : F(A|X_i) \rightarrow F(A|Y_i))_{(A,i) \in \mathcal{A} \times \mathcal{B}}$$

be the induced family in $\mathbf{Mnd}_{\mathcal{A}}(\mathcal{E})$. By definition, a morphism in $\mathbf{Mnd}_{\mathcal{A}}(\mathcal{E})$ is $F(\mathcal{A}|\mathcal{B})$ - λ -cellular for an ordinal λ if it is a λ -composite of pushouts of coproducts of maps in $F(\mathcal{A}|\mathcal{B})$; an object is λ -cellular if the unique map to it from the initial monad $\text{Id}_{\mathcal{E}}$ is λ -cellular. Accordingly, on taking \mathcal{V} -categories of algebras, a λ -cellular monad gives rise to a λ^+ -cochain of \mathcal{V} -categories and \mathcal{V} -functors, with base $\text{Alg}(\text{Id}_{\mathcal{E}}) \cong \mathcal{E}$.

Theorem 6.3. *Let $\mathcal{A} \hookrightarrow \mathcal{E}$ be a small dense full subcategory of a locally presentable category and \mathcal{B} a family of morphisms of \mathcal{E} . Given an ordinal λ , towers of $(\mathcal{A}, \mathcal{B})$ - λ -injectives coincide, up to isomorphism of λ^+ -cochains, with towers of algebras of $F(\mathcal{A}|\mathcal{B})$ - λ -cellular monads.*

Proof. Given Proposition 6.2 the proof follows inductively, arguing as in 3.12. □

Examples 6.4. Consider again $\mathcal{E} = \mathcal{V} = \mathbf{Cat}$, and $\mathbb{F} \hookrightarrow \mathbf{Cat}$ the dense full sub-2-category of finite cardinals, viewed as finite discrete categories and let

$$\mathcal{B} = \{j_0 : \emptyset \rightarrow (\bullet), j_1 : (\bullet \bullet) \rightarrow (\bullet \rightarrow \bullet), j_2 : (\bullet \rightrightarrows \bullet) \rightarrow (\bullet \rightarrow \bullet)\}$$

be the generating cofibrations for the folk model structure on \mathbf{Cat} [16]. Then structures such as monoidal categories — whose definition involves *no equations between objects* — can be described as $(\mathbb{F}, \mathcal{B})$ -iterated algebraic injectives in \mathbf{Cat} .

For a simple case, building on Example 3.11, let us describe categories C equipped with a tensor product $m : C^2 \rightarrow C : (x, y) \mapsto xy$ and an associator $\alpha : (xy)z \rightarrow x(yz)$ satisfying MacLane’s pentagon equation as 3-injectives in \mathbf{Cat} . From Example 6.1 objects of $\mathbb{I}nj(i_0)$ are categories equipped with a tensor product. The terms $t = (xy)z, s = x(yz)$ in three variables are elements of $UF3$ and, together, correspond to a single map $\langle t, s \rangle : F_1(\bullet \bullet) \rightarrow F_13$. Forming the pushout as in the second square below, we see that to equip (C, m) with the structure of an object of $\mathbb{I}nj(i_1)$ is to give a natural transformation $\alpha : (xy)z \rightarrow x(yz)$.

$$\begin{array}{ccccc} \emptyset & \longrightarrow & 2 & & F_1(\bullet \bullet) & \xrightarrow{(t,s)} & F_13 & & F_2(\bullet \rightrightarrows \bullet) & \xrightarrow{(u,v:l \rightrightarrows r)} & F_24 \\ j_0 \downarrow & & \downarrow i_0 & & F_1 j_1 \downarrow & & \downarrow i_1 & & F_2 j_2 \downarrow & & \downarrow i_2 \\ (\bullet) & \longrightarrow & 3 & & F_1(\bullet \rightarrow \bullet) & \longrightarrow & F_13 / \langle t \rightarrow s \rangle & & F_2(\bullet \rightarrow \bullet) & \longrightarrow & F_24 / \langle u = v \rangle \end{array}$$

Now let $l = ((wx)y)z$ and $r = w(x(yz))$. The two paths $u, v : l \rightrightarrows r$ of the pentagon specify a parallel pair in U_2F_24 and so correspond to a single map $F_2(\bullet \rightrightarrows \bullet) \rightarrow F_24$. Forming the pushout in the third square below, we see that (C, m, α) admits the structure of an object of $\mathbb{I}nj(i_2)$ precisely if the pentagon equation for α holds.

Building on this example, we can use $(\mathbb{F}, \mathcal{B})$ -3-injectivity to capture any monadic algebraic structure borne by categories, whose basic operations are of the form $C^n \rightarrow C$ for finite cardinals n , and whose defining equations only involving equalities between morphisms rather than objects. A precise characterisation of the 2-monads describing such categorical structures was given in Section 6 of [9], where they were called the *pie presentable* strongly finitary 2-monads. Indeed using Theorem 34 of *ibid*, Theorem 6.3 above and the pseudo-variant of Theorem 17 of [10], one can show that $(\mathbb{F}, \mathcal{B})$ - λ -injectives are $(\mathbb{F}, \mathcal{B})$ -3-injectives for all higher ordinals λ , with both classes coinciding as exactly the algebras for pie presentable strongly finitary 2-monads.

As mentioned already, Theorem 6.3 can be reformulated using \mathcal{A} -theories rather than monads. By Theorem 17 of [11] we have an equivalence of categories $\Psi : \mathbf{Th}_{\mathcal{A}}(\mathcal{E}) \cong \mathbf{Mnd}_{\mathcal{A}}(\mathcal{E})$. Translating through this equivalence we obtain a forgetful functor $U : \mathbf{Th}_{\mathcal{A}}(\mathcal{E}) \rightarrow \mathbf{Sig}_{\mathcal{A}}(\mathcal{E})$ with left adjoint F , abusively using the same symbols U and F as before. Since, by Theorem 24 of *ibid*, Ψ commutes with semantics in the sense that $\text{Mod}(\mathbb{T}) \simeq \text{Alg}(\Psi\mathbb{T})$, it follows that towers of algebras of $F(\mathcal{A}|\mathcal{B})$ - λ -cellular monads coincide, to within equivalence, with towers of models of $F(\mathcal{A}|\mathcal{B})$ - λ -cellular theories. Thus we may reformulate Theorem 6.3 using the language of \mathcal{A} -theories, as follows.

Theorem 6.5. *Let $\mathcal{A} \hookrightarrow \mathcal{E}$ be a small dense full subcategory of a locally presentable category and \mathcal{B} a family of morphisms of \mathcal{E} . Given an ordinal λ , towers of $(\mathcal{A}, \mathcal{B})$ - λ -injectives coincide, up to equivalence of λ -cochains, with towers of models of $F(\mathcal{A}|\mathcal{B})$ - λ -cellular theories.*

Example 6.6. In Section 3.11 of [19] Maltiniotis shows that the cellular globular theories, in the sense of Section 2.4, are exactly the ω -cellular theories relative to a set of maps

$$\mathcal{I} = \{ \Theta_0[\vec{m}, n] \rightarrow \Theta_0[\vec{m}, n]' \in \Theta_0\text{-Th} : \vec{m} \in \Theta_0, n \in \mathbb{N} \} .$$

These globular theories are characterised by universal properties — a morphism $\Theta_0[\vec{m}, n] \rightarrow \mathbb{T}$ amounts to a parallel pair $(n) \rightrightarrows \vec{m}$ in \mathbb{T} , whilst a morphism from $\Theta_0[\vec{m}, n]'$ to \mathbb{T} amounts to a morphism $(n + 1) \rightarrow \vec{m}$. Furthermore $\Theta_0[\vec{m}, n] \rightarrow \Theta_0[\vec{m}, n]'$ is the evident map determined by these universal properties.

To explain this using the framework of the present section, we work in ordinary **Set**-enriched category theory, with base $\mathcal{E} = [\mathbb{G}^{op}, \mathbf{Set}]$ and $\mathcal{A} = \Theta_0$. Then the Θ_0 -theories are — by Examples 42(v) of [11] — precisely the globular theories. Moreover, the forgetful functor $U : \Theta_0\text{-Th} \rightarrow [ob\Theta_0, [\mathbb{G}^{op}, \mathbf{Set}]]$ to signatures — to within isomorphism — sends $J : \Theta_0 \rightarrow \mathbb{T}$ to the signature $\vec{m} \mapsto T(JD-, \vec{m})$.

Now consider the set $\mathcal{B} = \{j_n : S(n) \rightarrow Y(n + 1) : n \in \mathbb{N}\}$ of boundary inclusions. To give a morphism of globular theories $F(\vec{m}|S(n)) \rightarrow \mathbb{T}$ is to give a morphism of signatures $\vec{m}|S(n) \rightarrow UT$. In turn, this amounts to a morphism $S(n) \rightarrow T(JD-, \vec{m})$; that is, a parallel pair $(n) \rightrightarrows \vec{m} \in T$. Therefore $F(\vec{m}|S(n)) \cong \Theta_0[\vec{m}, n]$. Continuing in this way we see, indeed, that the set \mathcal{I} considered by Maltiniotis is nothing but our canonical set of morphisms $F(\Theta_0|\mathcal{B})$. Given this identification, Theorem 3.12 becomes a special case of Theorem 6.5 above.

Appendix A: Fibrant cochains

In the present section we prove the results about fibrant cochains needed within the paper.

Proposition A.1. *Towers of models of cellular globular theories and of λ -injectives are smooth and isofibrant.*

Proof. Smoothness in the first case was observed in Section 2.4, whilst in the second case is by definition. Since isofibrations are defined by a right lifting property, they are stable under transfinite cocomposition. Given this, a smooth cochain X is isofibrant just when each link map $X_{n+1} \rightarrow X_n$ is an isofibration. Both classes of tower being smooth, it suffices to show that if $J : \mathbb{T} \rightarrow \mathbb{S}$ is a morphism of globular theories then $J^* : \text{Mod}(\mathbb{S}) \rightarrow \text{Mod}(\mathbb{T})$ and likewise that each forgetful functor of the form $V : \text{Inj}(\mathcal{I}) \rightarrow \mathcal{C}$ is an isofibration.

Since the structure of an algebraically injective object can be uniquely transferred along an isomorphism, the latter case is clear. As for the former, since each category of models is a *replete* full subcategory of the corresponding presheaf category, it suffices to show that $[J^{op}, 1] : [\mathbb{S}^{op}, \mathbf{Set}] \rightarrow [\mathbb{T}^{op}, \mathbf{Set}]$ is an isofibration. This follows easily from the fact that J is bijective on objects. □

The following results about the good properties of smooth isofibrant cochains are naturally 2-categorical — as such, we state and prove them using general 2-categories rather than just *CAT*. Here a 2-category refers to a *strict* 2-category. Given 2-categories \mathcal{A} and \mathcal{B} the collection of strict 2-functors, pseudonatural transformations and modifications between them forms a 2-category $Ps(\mathcal{A}, \mathcal{B})$. Limits and isofibrations are representable notions in a 2-category — thus,

a cochain $(X_n)_{n \in \lambda} \in \mathcal{C}$ is said to be smooth/isofibrant if for each $C \in \mathcal{C}$ the induced cochain $\mathcal{C}(C, X_n)_{n \in \lambda}$ of categories is smooth/isofibrant.

Lemma A.2. *Consider a 2-category \mathcal{C} and a pseudonatural transformation $f : X \rightsquigarrow Y$ of λ -cochains in \mathcal{C} . If Y is smooth and isofibrant, then there exists a 2-natural transformation $g : X \rightarrow Y$ and invertible modification $\phi : f \cong g$.*

Proof. If $\lambda = 0$ then it is trivial. For λ non-zero, we define $g_0 = f_0 : X_0 \rightarrow Y_0$ and $\theta_0 = id$. For the inductive step suppose that $\alpha < \lambda$ and that we have defined $g_n : X_n \rightarrow Y_n$ for $n < \alpha$ naturally in n , as well as invertible 2-cells $\phi_n : f_n \cong g_n$ satisfying the modification equation at $m < n < \alpha$. In summary, we have an invertible modification $\theta_{<\alpha} : f_{<\alpha} \cong g_{<\alpha}$. To prove the result, by transfinite induction, we must extend it to $\theta_{\leq\alpha} : f_{\leq\alpha} \cong g_{\leq\alpha}$.

If $\alpha = \beta + 1$ we have the diagram below left.

$$\begin{array}{ccc}
 \begin{array}{ccc}
 X_{\beta+1} & \xrightarrow{f_{\beta+1}} & Y_{\beta+1} \\
 \downarrow x_{\beta}^{\beta+1} & \searrow f_{\beta}^{\beta+1} \Downarrow & \downarrow y_{\beta}^{\beta+1} \\
 X_{\beta} & \xrightarrow{f_{\beta}} & Y_{\beta} \\
 & \searrow \theta_{\beta} \Downarrow & \\
 & & Y_{\beta} \\
 & \xrightarrow{g_{\beta}} &
 \end{array} & = &
 \begin{array}{ccc}
 X_{\beta+1} & \xrightarrow{f_{\beta+1}} & Y_{\beta+1} \\
 \downarrow x_{\beta}^{\beta+1} & \searrow \theta_{\beta+1} \Downarrow & \downarrow y_{\beta}^{\beta+1} \\
 X_{\beta} & \xrightarrow{g_{\beta}} & Y_{\beta} \\
 & \searrow \parallel & \\
 & & Y_{\beta} \\
 & \xrightarrow{g_{\beta}} &
 \end{array}
 \end{array}$$

Since $\mathcal{C}(X_{\beta+1}, y_{\beta}^{\beta+1}) : \mathcal{C}(X_{\beta+1}, Y_{\beta+1}) \rightarrow \mathcal{C}(X_{\beta+1}, Y_{\beta})$ is an isofibration there exists $\theta_{\beta+1}$ and $g_{\beta+1}$ as on the right, making the two diagrams equal. The equation is precisely the modification condition at $\beta < \beta + 1$ so that we have $g_{\leq\beta+1} : X_{\leq\beta+1} \rightarrow Y_{\leq\beta+1}$ and $\theta_{\leq\beta+1} : f_{\leq\beta+1} \cong g_{\leq\beta+1}$, as required.

If α is a limit ordinal then since $Y_{\alpha} = \lim_{n < \alpha} Y_n$ we obtain a unique morphism $g_{\alpha} : X_{\alpha} \rightarrow Y_{\alpha}$ satisfying $y_n^{\alpha} \circ g_{\alpha} = g_n \circ y_n^{\alpha}$ for each $n < \alpha$. This gives the desired cochain map $g_{\leq\alpha} : X_{\leq\alpha} \rightarrow Y_{\leq\alpha}$. It remains to provide an invertible 2-cell $\theta_{\alpha} : f_{\alpha} \cong g_{\alpha}$ satisfying the modification condition


$$\begin{array}{ccc}
 \begin{array}{ccc}
 X_{\alpha} & \xrightarrow{f_{\alpha}} & Y_{\alpha} \\
 \downarrow x_n^{\alpha} & \searrow \theta_{\alpha} \Downarrow & \downarrow y_n^{\alpha} \\
 X_n & \xrightarrow{g_n} & Y_n \\
 & \searrow \parallel & \\
 & & Y_n \\
 & \xrightarrow{g_n} &
 \end{array} & = &
 \begin{array}{ccc}
 X_{\alpha} & \xrightarrow{f_{\alpha}} & Y_{\alpha} \\
 \downarrow x_n^{\alpha} & \searrow f_n^{\alpha} \Downarrow & \downarrow y_n^{\alpha} \\
 X_n & \xrightarrow{f_n} & Y_n \\
 & \searrow \theta_n \Downarrow & \\
 & & Y_n \\
 & \xrightarrow{g_n} &
 \end{array}
 \end{array} \tag{8}$$

for each $n < \alpha$. Now the composite 2-cells in the diagram above right are of the form

$$\phi_n : y_n^{\alpha} \circ f_{\alpha} \Rightarrow g_n \circ x_n^{\alpha} = y_n^{\alpha} \circ g_{\alpha}$$

and indeed satisfy $y_m^{\alpha} \circ \phi_n = \phi_m$ for all $m < n$. This is proven by the following equations of

2-cells



of which the first uses that $\theta : f_{<\alpha} \Rightarrow g_{<\alpha}$ is a modification and the second uses pseudonaturality of f . By the universal property of the limit $\mathcal{C}(X_\alpha, Y_\alpha)$ in **CAT** therefore there exists a unique 2-cell θ_α which, on postcomposition by the y_n^α equals ϕ_n ; in other words, satisfying the required equation (8) for a modification depicted above. \square

- Proposition A.3.** 1. If $X : \lambda^{op} \rightarrow \mathcal{C}$ is a smooth isofibrant cochain then at each limit ordinal $\alpha < \lambda$ the limit $X_\alpha = \lim_{n < \alpha} X_n$ is a bilimit.
 2. Each pseudonatural equivalence between smooth isofibrant λ -cochains is isomorphic (via a modification) to a cochain equivalence. In particular, cochain equivalence is an equivalence relation, when restricted to smooth isofibrant cochains.

Proof. 1. To say that the cone $\Delta(X_\alpha) \rightarrow X_{<\alpha} \in [\alpha^{op}, \mathcal{C}]$ exhibits the limit X_α as a bilimit is, by definition, to say that the composite

$$\mathcal{C}(A, X_\alpha) \rightarrow [\alpha^{op}, \mathcal{C}](\Delta A, X_{<\alpha}) \hookrightarrow Ps(\alpha^{op}, \mathcal{C})(\Delta A, X_{<\alpha})$$

is an equivalence for each A . Since the first component is an isomorphism, this will be the case just when the second component is an equivalence. Since it is always fully faithful, this is will be the case just when it is essentially surjective on objects and, by the preceding lemma, this holds since $X_{<\alpha}$ is smooth isofibrant.

2. If $f : X \rightsquigarrow Y$ is a pseudonatural equivalence, then by the lemma there is a 2-natural transformation $g : X \rightarrow Y$ and invertible modification $f \cong g$. For $n \in \lambda$ we have $f_n \cong g_n$ since each f_n is an equivalence so too is the naturally isomorphic g_n . This proves the first part. The second part from the first part using that pseudonatural equivalence is an equivalence relation. \square

Acknowledgements

The main results of this paper were presented at CT2016 in Halifax [7]. I am grateful to the organisers for inviting me to give a talk, and for the very enjoyable conference. I am also grateful to Dimitri Ara and Georges Maltsiniotis for encouraging me to publish the results described herein. I acknowledge the support of the Grant Agency of the Czech Republic under the grant 19-00902S.

References

- [1] ADÁMEK, J. AND ROSICKÝ, J. *Locally Presentable and Accessible Categories*, vol. 189 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, 1994.
- [2] ARA, DIMITRI. Sur les ∞ -groupoïdes de Grothendieck et une variante ∞ -catégorique. PhD thesis, Université Paris VII, 2010.
- [3] ARA, DIMITRI. On the homotopy theory of Grothendieck ω -groupoids. *Journal of Pure and Applied Algebra* 217 (2013), 1237–1278.
- [4] BATANIN, MICHAEL. Monoidal globular categories as a natural environment for the theory of weak n -categories. *Advances in Mathematics* 136(1) (1998) 39–103.
- [5] BERGER, CLEMENS. A Cellular Nerve for Higher Categories. *Advances in Mathematics* 169,(2002) 118–175.
- [6] BIRD, GREGORY. *Limits in 2-categories of locally-presented categories*. PhD thesis, University of Sydney, 1984.
- [7] BOURKE, JOHN. Grothendieck ω -groupoids as iterated injectives – talk given at CT2016. Slides at <http://mysite.science.uottawa.ca/phofstra/CT2016/slides/Bourke.pdf>.
- [8] BOURKE, JOHN. Equipping weak equivalences with algebraic structure. *Mathematische Zeitschrift* 294(3) (2020), 995-1019.
- [9] BOURKE, J. AND GARNER, R. On semiflexible, flexible and pie algebras. *Journal of Pure and Applied Algebra* 217 (2013), 293–321.
- [10] BOURKE, J. AND GARNER, R. Algebraic weak factorisation systems II: Categories of weak maps. *Journal of Pure and Applied Algebra* 220 (2016), 148–174.
- [11] BOURKE, J. AND GARNER, R. Monads and theories. *Advances in Mathematics* 251 (2019), 1024–1071.
- [12] GABRIEL, P., AND ULMER, F. *Lokal präsentierbare Kategorien*, vol. 221 of *Lecture Notes in Mathematics*. Springer-Verlag, 1971.
- [13] GARNER, RICHARD. Understanding the small object argument. *Applied Categorical Structures* 17(3) (2009), 247–285.
- [14] GROTHENDIECK, ALEXANDER. *Pursuing Stacks* (1983).
- [15] HENRY, SIMON. Algebraic models of homotopy types and the homotopy hypothesis. Arxiv Preprint 2016, <https://arxiv.org/abs/1609.04622>.
- [16] LACK, STEVE. Homotopy-theoretic aspects of 2-monads. *Journal of Homotopy and Related Structures* 7(2) (2007), 229–260.
- [17] LANARI, EDOARDO. Towards a globular path object for weak ∞ -groupoids. *Journal of Pure and Applied Algebra* 224 (2020), 630–702.
- [18] LEINSTER, TOM. A survey of definitions of n -category. *Theory and Applications of Categories* 10 (2002) 1–70.

- [19] MALTSINIOTIS, GEORGES. Grothendieck ω -groupoids, and still another definition of ω -categories. Arxiv Preprint 2010, <https://arxiv.org/abs/1009.2331>.
- [20] NIKOLAUS, THOMAS. Algebraic models for higher categories. *Indag. Math. (N.S.)* 21 (2011), no. 1–2, 52–75.
- [21] STREET, ROSS. The petit topoi of globular sets. *Journal of Pure and Applied Algebra* 154 (2000) 299–315.