

Unitary dual functors for unitary multitensor categories

David Penneys^a

^aThe Ohio State University, Columbus OH 43210, USA

Abstract

We classify which dual functors on a unitary multitensor category are compatible with the dagger structure in terms of groupoid homomorphisms from the universal grading groupoid to $\mathbb{R}_{>0}$ where the latter is considered as a groupoid with one object. We then prove that all unitary dual functors induce unitarily equivalent bi-involutive structures. As an application, we provide the unitary version of the folklore correspondence between shaded planar C^* algebras with finite dimensional box spaces and unitary multitensor categories with a chosen unitary dual functor and chosen generator. We make connection with the recent work of Giorgetti-Longo to determine when the loop parameters in these planar algebras are scalars. Finally, we show that we can correct for many non-spherical choices of dual functor by adding the data of a spherical state on $\text{End}_{\mathcal{C}}(1_{\mathcal{C}})$, similar to the spherical state for a graph planar algebra.

This is the published version of [arXiv:1808.00323](https://arxiv.org/abs/1808.00323).

Received: 2nd August, 2018. Accepted: 4th May, 2020.

MSC: 18D05 and 18D10 primary, 46L37 secondary.

Keywords: tensor category, planar algebra, pivotal structure, unitary dual functor.

1. Introduction

In a rigid monoidal category \mathcal{C} , every object has a *dual*, consisting of a triple $(c^{\vee}, \text{ev}_c, \text{coev}_c)$ where $c^{\vee} \in \mathcal{C}$ and $\text{ev}_c \in \mathcal{C}(c^{\vee} \otimes c \rightarrow 1_{\mathcal{C}})$ and $\text{coev}_c \in \mathcal{C}(1_{\mathcal{C}} \rightarrow c \otimes c^{\vee})$ satisfy the *zig-zag axioms*:

$$\begin{array}{c} c \\ \lrcorner \\ \text{---} \\ \lrcorner \\ c^{\vee} \\ \lrcorner \\ \text{---} \\ \lrcorner \\ c \end{array} := (\text{id}_c \otimes \text{ev}_c) \circ (\text{coev}_c \otimes \text{id}_c) = \text{id}_c =: \begin{array}{c} \lrcorner \\ \text{---} \\ \lrcorner \\ c \end{array} \qquad \begin{array}{c} \lrcorner \\ \text{---} \\ \lrcorner \\ c^{\vee} \\ \lrcorner \\ \text{---} \\ \lrcorner \\ c \end{array} = \begin{array}{c} \lrcorner \\ \text{---} \\ \lrcorner \\ c^{\vee} \end{array} = \text{id}_{c^{\vee}},$$

and every object is isomorphic to the dual of some other object. By choosing a dual for each $c \in \mathcal{C}$, we get an anti-monoidal *dual functor* $\vee : \mathcal{C} \rightarrow \mathcal{C}$ defined on a morphism $f \in \mathcal{C}(a \rightarrow b)$ by

$$f^{\vee} := \begin{array}{c} \lrcorner \\ \text{---} \\ \lrcorner \\ a^{\vee} \\ \lrcorner \\ \text{---} \\ \lrcorner \\ b^{\vee} \end{array} = (\text{ev}_d \otimes \text{id}_{c^{\vee}}) \circ (\text{id}_{d^{\vee}} \otimes f \otimes \text{id}_{c^{\vee}}) \circ (\text{id}_{d^{\vee}} \otimes \text{coev}_c).$$

Email address: penneys.2@osu.edu (David Penneys)

© David Penneys, 2020, under a Creative Commons Attribution 4.0 International License.

DOI: [10.21136/HS.2020.09](https://doi.org/10.21136/HS.2020.09)

A dual functor comes with a canonical anti-monoidal tensorator

$$\begin{aligned} \nu_{a,b} &:= \text{diagram: a cup with a vertical line on the left and a vertical line on the right, with an arrow labeled $(a \otimes b)^\vee$ pointing down from the right side. The left side is labeled b^\vee and the right side is labeled a^\vee} \\ &= (\text{ev}_b \otimes \text{id}_{(a \otimes b)^\vee}) \circ (\text{id}_{b^\vee} \otimes \text{ev}_a \otimes \text{id}_b \otimes \text{id}_{(a \otimes b)^\vee}) \circ (\text{id}_{b^\vee \otimes a^\vee} \otimes \text{coev}_{a \otimes b}), \end{aligned}$$

and any two dual functors \vee_1, \vee_2 are *uniquely* monoidally naturally isomorphic via

$$\zeta_c := \text{diagram: a cup with a vertical line on the left and a vertical line on the right, with an arrow labeled c^{\vee_1} pointing down from the right side. The left side is labeled c^{\vee_2}} = (\text{ev}_c^2 \otimes \text{id}_{c^{\vee_1}}) \circ (\text{id}_{c^{\vee_2}} \otimes \text{coev}_c^1).$$

A *pivotal structure* on \mathcal{C} is a pair (\vee, φ) , where \vee is a chosen dual functor, and $\varphi : \text{id} \Rightarrow \vee \circ \vee$ is a monoidal natural isomorphism. Using φ , one can define the left and right *quantum dimension* of an object $c \in \mathcal{C}$; we refer the reader to §2 for a detailed discussion of pivotal structures.

The above definitions are poorly behaved in the context of rigid tensor C^* categories. More precisely, the above definitions fail the *principle of equivalence* [nLa18], which roughly states that mathematical definitions should be invariant under the proper notion of equivalence. As a basic example, when one works with Hilbert spaces, the correct notion of equivalence is that of *unitary isomorphism*, i.e., bounded linear maps $u : H \rightarrow K$ such that $u^*u = \text{id}_H$ and $uu^* = \text{id}_K$, and *not* bounded linear isomorphism.¹ As C^* categories admit an equivalent definition as those categories which admit a faithful dagger functor to the category \mathbf{Hilb} of Hilbert spaces which is norm-closed on the level of hom spaces [GLR85], we see that one must work with dagger functors and unitary (natural) isomorphisms to satisfy the principle of equivalence for dagger categories.

Indeed, a dual functor on a rigid tensor C^* category need not be a dagger functor, and the canonical tensorator ν need not be unitary. With this in mind, we call a dual functor $\vee : \mathcal{C} \rightarrow \mathcal{C}$ *unitary* if it is a dagger tensor functor, i.e., for all $a, b \in \mathcal{C}$ and $f \in \mathcal{C}(a \rightarrow b)$, the canonical tensorator $\nu_{a,b}$ is unitary and $f^{\vee\dagger} = f^{\vee}$. Given a unitary dual functor $\vee : \mathcal{C} \rightarrow \mathcal{C}^{\text{mop}}$, there is a unique pivotal structure for which left and right dimensions of objects are positive² and given by $\text{ev}_c^\dagger \circ \text{ev}_c$ and $\text{coev}_c \circ \text{coev}_c^\dagger$ respectively:

$$\varphi_c := (\text{coev}_c^\dagger \otimes \text{id}_{c^{\vee\vee}}) \circ (\text{id}_c \otimes \text{coev}_{c^\vee}) = (\text{id}_{c^{\vee\vee}} \otimes \text{ev}_c) \circ (\text{ev}_{c^\vee}^\dagger \otimes \text{id}_c). \quad (1)$$

By [Sel11, Lem. 7.5] (which is Proposition 3.9 below), a dual functor \vee is unitary if and only if φ defined as in (1) above defines a pivotal structure. We call such pivotal structures *unitary*, but one should really only consider the term ‘unitary pivotal structure’ as a synonym for ‘the canonical pivotal structure associated to a unitary dual functor’ as in [Sel11, §7.3].

Unitary dual functors on rigid tensor C^* categories were first constructed in [LR97, Yam04, BDH14]. The notion of a quantum dimension for dualizable objects in a tensor C^* category with simple unit object was established in [LR97] via *standard solutions to the conjugate equations*. In [Yam04], it was further clarified that for a *unitary tensor category* \mathcal{C} , which is an idempotent

¹Conjugating a self-adjoint operator by a bounded linear isomorphism need not produce a self-adjoint operator unless the isomorphism is unitary. Similarly, in finite dimensions, taking coordinates for a self-adjoint operator with respect to a basis need not produce a self-adjoint matrix unless the basis is orthonormal. In this respect, the notions of linear isomorphism and basis fail the principle of equivalence for the C^* category $\mathbf{Hilb}_{\text{fd}}$ of finite dimensional Hilbert spaces, while the notions of unitary isomorphism and orthonormal basis satisfy the principle of equivalence.

²We call a pivotal structure *pseudounitary* if all quantum dimensions of objects are strictly positive. This definition is equivalent to [EGNO15, Def. 9.4.4] for fusion categories by uniqueness of the Frobenius-Perron dimensions.

complete rigid tensor C^* category with simple unit object, for every object $c \in \mathcal{C}$, there is a unique *balanced* dual $(\bar{c}, \text{ev}_c, \text{coev}_c)$ up to unique unitary isomorphism satisfying the zig-zag axioms and the balancing equation:

$$\text{ev}_c \circ (\text{id}_{\bar{c}} \otimes f) \circ \text{ev}_c^\dagger = \text{coev}_c^\dagger \circ (f \otimes \text{id}_{\bar{c}}) \circ \text{coev}_c \quad \forall c \in \mathcal{C}, f \in \mathcal{C}(c \rightarrow c).$$

Moreover, choosing these balanced duals gives a canonical unitary dual functor whose associated unitary pivotal structure is *spherical*. This result was later expanded in [BDH14], in the context of von Neumann algebras with finite dimensional centers, to *unitary multitensor categories*, which are idempotent complete rigid tensor C^* categories. For a unitary multitensor category \mathcal{C} , $1_{\mathcal{C}}$ is no longer simple; however, since \mathcal{C} is automatically semisimple by a generalization of [LR97, Lem. 3.9], $1_{\mathcal{C}}$ decomposes as an orthogonal finite direct sum of simples $1_{\mathcal{C}} = \bigoplus_{i=1}^r 1_i$. Each ‘corner’ $\mathcal{C}_{ii} := 1_i \otimes \mathcal{C} \otimes 1_i$ is again a unitary tensor category.

While the existence of this canonical unitary dual functor and spherical structure for a unitary multitensor category is extremely powerful, it is not always the most relevant unitary dual functor for applications. A first example is the unitary tensor category $\text{Bim}_{\text{bf}}(R)$ of bifinite bimodules over the hyperfinite II_1 factor R . The most widely used unitary dual functor on $\text{Bim}_{\text{bf}}(R)$ is built from the canonical trace on R via left and right R -valued inner products on the subspaces of bounded vectors (see [Bis97, Pen13, AP17, JP19]). Often, one restricts to *spherical/extremal* bimodules where the canonical unitary dual functor agrees with this tracial one.

Notice $\text{Bim}_{\text{bf}}(R)$ admits a grading by $\mathbb{R}_{>0}$ given by the ratio of left to right von Neumann dimension. Whether $\mathbb{R}_{>0}$ is the universal grading group of $\text{Bim}_{\text{bf}}(R)$ is a tantalizing open question. However, this grading is sufficient to understand the difference between the tracial unitary dual functor and unitary pivotal structure, which corresponds to the identity group homomorphism $\mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$, and the canonical unitary spherical structure, which corresponds to the trivial group homomorphism under our Theorem A below. We refer the reader to Example 3.43 for more details.

A second example is the industry of constructing subfactor planar algebras as planar subalgebras of graph planar algebras [Jon01, Gup08, Pet10, BMPS12, Han10, MP15a, MP15b, PP15, LMP15, GMP⁺18]. (Such a realization is always possible for finite depth subfactor planar algebras by [JP11], although this result is not necessary in the construction. See also [GMP⁺18, CHPS18] for the module embedding theorem.) By Example 4.7 below, the projection category of the planar algebra of a finite connected bipartite graph Γ is dagger equivalent to $\text{End}^\dagger(\text{Hilb}_{\text{fd}}^n)$, the unitary multitensor category of dagger endofunctors of n copies of finite dimensional Hilbert spaces, where n is the number of vertices of Γ . The planar algebra gives a particular unitary pivotal structure related to Frobenius-Perron data of Γ , which does *not* correspond to the canonical unitary spherical structure. However, one has a canonical *spherical state* on the graph planar algebra [Jon00, Prop. 3.4], which implies any *evaluatable* planar subalgebra is a subfactor planar algebra [Jon01, §8]. We refer the reader to §5 for more details.

The relevant unitary dual functor and corresponding pivotal structure to explain this second class of examples is provided by [GL19] in the context of 2- C^* -categories, which defines standard/minimal solutions to the conjugate equations *with respect to a particular object* $X \in \mathcal{C}$. While providing deeper insight into well-behaved choices of solutions to the conjugate equations, they leave open the important question of classifying all unitary dual functors. Notice that although two unitary dual functors are uniquely monoidally naturally isomorphic, this natural isomorphism need not be unitary! (The unique monoidal natural isomorphism may fail the

principle of equivalence for tensor C^* categories.) Hence two unitary dual functors need not be unitarily equivalent.

In this article, we prove the following classification theorem.

Theorem A. *Let \mathcal{C} be a unitary multitensor category. There are canonical bijections between:*

- (1) *Pseudounitary³ pivotal structures up to monoidal natural isomorphism.⁴*
- (2) *Unitary dual functors up to unitary monoidal natural isomorphism.⁵*
- (3) *Groupoid homomorphisms $\mathcal{U} \rightarrow \mathbb{R}_{>0}$, where the latter is viewed as a groupoid with one object.*

Here, \mathcal{U} is the *universal grading groupoid* of \mathcal{C} , which is defined analogously to the universal grading group of a tensor category as in [EGNO15, §4.14] (see §3.3 below for the definition). We show in Lemma 3.26 that from a pseudounitary pivotal structure φ , we get a groupoid homomorphism by taking ratios of dimensions of simple objects; that is, if a simple $c \in \mathcal{C}$ is graded by a morphism $g \in \mathcal{U}$, we get a well defined $\pi \in \text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$ by

$$\pi(g) := \frac{\dim_L^\varphi(c)}{\dim_R^\varphi(c)}.$$

Conversely, following a suggestion of André Henriques, given a $\pi \in \text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$, we define a canonical π -balanced dual functor \vee_π which is unique up to unique unitary monoidal natural isomorphism by finding π -balanced solutions $(\text{ev}_c^\pi, \text{coev}_c^\pi)$ to the conjugate equations which satisfy the zig-zag axioms and for all $f \in \mathcal{C}(c \rightarrow c)$ and morphisms $g \in \mathcal{U}$,

$$\Psi \left(\text{ev}_c^\pi \circ (\text{id}_{c \vee_\pi} \otimes f_g) \circ (\text{ev}_c^\pi)^\dagger \right) = \pi(g) \cdot \Psi \left((\text{coev}_c^\pi)^\dagger \circ (f_g \otimes \text{id}_{c \vee_\pi}) \circ \text{coev}_c^\pi \right)$$

where Ψ the linear functional in $\mathcal{C}(1 \rightarrow 1) \rightarrow \mathbb{C}$ which sends every minimal projection to $1_{\mathbb{C}}$, $f_g \in \mathcal{C}(c_g \rightarrow c_g)$ is the g -homogeneous component of f , and c_g is the g -graded component of $c \in \mathcal{C}$. This proof is similar to [Yam04, Lem. 3.9] and [NT13, Prop. 2.2.15].

Unitary dual functors and pivotal structures are closely related to the more general notion of *bi-involutive structure* from [HP17, §2.1]. An *involution* on a multitensor category [Egg11] is a conjugate-linear anti-monoidal functor $(\bar{\cdot}, \nu) : \mathcal{C} \rightarrow \mathcal{C}$ together with a monoidal natural isomorphism $\varphi : \text{id}_{\mathcal{C}} \Rightarrow \bar{\bar{\cdot}}$. When \mathcal{C} is unitary, we call $(\bar{\cdot}, \nu, \varphi)$ *bi-involutive* if $(\bar{\cdot}, \nu)$ is an anti-monoidal dagger functor and φ is unitary. One obtains a bi-involutive structure from a unitary dual functor \vee and its canonical unitary pivotal structure φ by simply forgetting the evaluation and coevaluation maps.

Motivated by the example $\text{Bim}_{\text{bf}}(R)$ above and [JP19, Rem. 2.14] (see also Example 3.43), we prove the following somewhat surprising result in §3.5.

Corollary B. *Any two bi-involutive structures on a unitary multitensor category induced by unitary dual functors are canonically unitarily equivalent.*

As an application of Theorem A, we now understand the unitary version of the folklore correspondence between shaded planar algebras and pivotal multitensor categories with a choice of generator [MPS10, Gho11, Yam12, BHP12, HPT16b].

³See Footnote 2 on the previous page for the definition of a pseudounitary pivotal structure.

⁴If two pseudounitary pivotal structures are monoidally naturally isomorphic, they are so in a unique way.

⁵If two unitary dual functors are unitarily monoidally naturally isomorphic, they are so in a unique way.

Theorem C. *There is an equivalence of categories⁶*

$$\left\{ \begin{array}{l} \text{Shaded planar } \mathbb{C}^* \text{ algebras} \\ \mathcal{P}_\bullet \text{ with finite dimensional} \\ \text{box spaces } \mathcal{P}_{n,\pm} \end{array} \right\} \cong \left\{ \begin{array}{l} \text{Triples } (\mathcal{C}, \vee, X) \text{ with } \mathcal{C} \text{ a unitary multitensor category,} \\ \vee \text{ a unitary dual functor, and a generator } X \in \mathcal{C} \text{ with} \\ \text{an orthogonal decomposition } 1_{\mathcal{C}} = 1_+ \oplus 1_- \text{ such that} \\ X = 1_+ \otimes X \otimes 1_- \end{array} \right\}$$

Here, we call $X \in \mathcal{C}$ a generator if every object of \mathcal{C} is isomorphic to a direct summand of alternating tensor powers of X and X^\vee .

The proof of this theorem is sketched in §4.1 below.

As mentioned earlier, for a chosen generator $X \in \mathcal{C}$ such that $1_{\mathcal{C}} = 1_+ \oplus 1_-$ and $X = 1_+ \otimes X \otimes 1_-$ as in Theorem C, the canonical standard/minimal solutions to the conjugate equations with respect to $X \in \mathcal{C}$ from [GL19] give a canonical unitary dual functor which makes both loop moduli identical scalars in the corresponding shaded planar \mathbb{C}^* algebra:

$$\text{○} = d_X \text{id}_{1_+} \qquad \text{○} = d_X \text{id}_{1_-} .$$

There is also a canonical unitary dual functor giving a unitary version of the *lopsided* convention from [MP14, §1.1] which has been instrumental for constructing many subfactor planar algebras as planar subalgebras of graph planar algebras. We refer the reader to §4.2 for more details.

As a final application, in §5, we ‘correct’ for some non-spherical choices of unitary pivotal structure on a unitary multitensor category \mathcal{C} . If \mathcal{C} is faithfully graded by \mathcal{M}_r , the groupoid with r objects and exactly one isomorphism between any two objects, then any groupoid homomorphism $\pi : \mathcal{M}_r \rightarrow \mathbb{R}_{>0}$ induces a unitary pivotal structure on \mathcal{C} by Theorem A and universality of \mathcal{U} . As usual, picking $\pi = 1$ gives the canonical unitary spherical structure.

Theorem D. *Suppose $\dim(\mathcal{C}(1_{\mathcal{C}} \rightarrow 1_{\mathcal{C}})) = r$ and \mathcal{C} is faithfully graded by \mathcal{M}_r . For each $\pi \in \text{Hom}(\mathcal{M}_r \rightarrow \mathbb{R}_{>0})$, there exists a unique faithful state $\psi^\pi : \mathcal{C}(1_{\mathcal{C}} \rightarrow 1_{\mathcal{C}}) \rightarrow \mathbb{C}$ such that for all $c \in \mathcal{C}$ and $f \in \mathcal{C}(c \rightarrow c)$, $\psi^\pi(\text{tr}_L^\pi(f)) = \psi^\pi(\text{tr}_R^\pi(f))$.*

This theorem generalizes the existence of the spherical state on the bipartite graph planar algebra from [Jon00, Prop. 3.4] which allows one to construct subfactor planar algebras by finding evaluable planar subalgebras. There is also a notion of a spherical state with respect to an object $X \in \mathcal{C}$ from [GL19, (7.9)]; we explain the relation between the two conventions in Example 5.11.

2. Pivotal structures

In what follows \mathcal{C} denotes a \mathbb{C} -linear category. We write $c \in \mathcal{C}$ to denote c is an object of \mathcal{C} and we write $\mathcal{C}(a \rightarrow b)$ for $\text{Hom}_{\mathcal{C}}(a, b)$.

Definition 2.1 ([EGNO15, Def. 4.1.1]). A multitensor category is a locally finite \mathbb{C} -linear abelian rigid monoidal category such that $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ is bilinear. We call \mathcal{C} *indecomposable* if it is not equivalent to the direct sum of two nonzero multitensor categories. If $\mathcal{C}(1_{\mathcal{C}} \rightarrow 1_{\mathcal{C}})$ is one-dimensional, i.e., $1_{\mathcal{C}}$ is *simple*, we call \mathcal{C} a tensor category.

We refer the reader to [EGNO15] for basic background material on multitensor categories. To ease the notation, whenever possible, we suppress the associator and unitor natural isomorphisms. All results on pivotal categories in §2.1 – 2.3 are well known to experts. We provide some proofs for completeness and convenience.

⁶Here we suppress a subtlety about contractible 2-categories; see Footnote 14 in Theorem 4.1 for details.

2.1 Pivotal categories We start by recalling the standard definition of a dual functor and a pivotal category. For this section, \mathcal{C} is a rigid monoidal category. This means for each $c \in \mathcal{C}$, there exists a dual object $c^\vee \in \mathcal{C}$ together with evaluation and coevaluation morphisms $\text{ev}_c, \text{coev}_c$ which satisfy the zig-zag axioms, and that for each $c \in \mathcal{C}$, there is a $c_\vee \in \mathcal{C}$ such that $(c_\vee)^\vee \cong c$.

Definition 2.2. A choice of dual $(c^\vee, \text{ev}_c, \text{coev}_c)$ for each $c \in \mathcal{C}$ assembles into a *dual functor*, which is a strong monoidal functor $\vee : \mathcal{C} \rightarrow \mathcal{C}^{\text{mop}}$ ⁷ defined on $f : \mathcal{C}(c \rightarrow d)$ by

$$f^\vee := \int_{d^\vee}^{\text{coev}_d} \left(\int_{c^\vee}^{\text{ev}_c} f \right) = (\text{ev}_d \otimes \text{id}_{c^\vee}) \circ (\text{id}_{d^\vee} \otimes f \otimes \text{id}_{c^\vee}) \circ (\text{id}_{d^\vee} \otimes \text{coev}_c).$$

A dual functor \vee comes with a canonical tensorator $\nu = \{\nu_{a,b} : a^\vee \otimes_{\mathcal{C}^{\text{mop}}} b^\vee := b^\vee \otimes a^\vee \rightarrow (a \otimes b)^\vee\}$ given by

$$\begin{aligned} \nu_{a,b} &:= \int_{a^\vee}^{\text{coev}_a} \int_{b^\vee}^{\text{ev}_b} (a \otimes b)^\vee \\ &= (\text{ev}_b \otimes \text{id}_{(a \otimes b)^\vee}) \circ (\text{id}_{b^\vee} \otimes \text{ev}_a \otimes \text{id}_b \otimes \text{id}_{(a \otimes b)^\vee}) \circ (\text{id}_{b^\vee} \otimes a^\vee \otimes \text{coev}_{a \otimes b}) \end{aligned} \quad (2)$$

and unit isomorphism $r := \text{coev}_{1_c} : 1 \rightarrow 1^\vee$. In what follows, we suppress r to ease the notation.

Remark 2.3. Given a dual $(c^\vee, \text{ev}_c, \text{coev}_c)$ of $c \in \mathcal{C}$, the morphism ev_c is completely determined by coev_c . (Similarly, ev_c completely determines coev_c .) Hence if $(c_i^\vee, \text{ev}_c^i, \text{coev}_c^i)$ for $i = 1, 2$ are two duals of c and if there is a $\zeta_c \in \mathcal{C}(c_2^\vee \rightarrow c_1^\vee)$ such that $(\text{id}_c \otimes \zeta_c) \circ \text{coev}_c^2 = \text{coev}_c^1$, then

$$\zeta_c = \int_{c_2^\vee}^{\text{coev}_{c_2}} \int_{c_1^\vee}^{\text{ev}_{c_1}} = (\text{ev}_c^2 \otimes \text{id}_{c_1^\vee}) \circ (\text{id}_{c_2^\vee} \otimes \text{coev}_c^1) \quad (3)$$

which is necessarily invertible, and $\text{ev}_c^1 \circ (\zeta_c \otimes \text{id}_c) = \text{ev}_c^2$ as well. Hence any two choices of dual functor are *uniquely* monoidally naturally isomorphic.

Moreover, given a dual functor \vee , the tensorator ν from (2) above is *not* part of the data of \vee , as it is the unique isomorphism $\zeta_{a \otimes b}$ for the two duals $((a \otimes b)^\vee, \text{ev}_{a \otimes b}, \text{coev}_{a \otimes b})$ and $(b^\vee \otimes a^\vee, \text{ev}_b \circ (\text{id}_{b^\vee} \otimes \text{ev}_a \otimes \text{id}_b), (\text{id}_a \otimes \text{coev}_b \otimes \text{id}_{a^\vee}) \circ \text{coev}_a)$.

Definition 2.4. A *pivotal structure* on a rigid monoidal category \mathcal{C} is a pair (\vee, φ) where $\vee : \mathcal{C} \rightarrow \mathcal{C}^{\text{mop}}$ is a dual functor and $\varphi : \text{id} \Rightarrow \vee \circ \vee$ is a monoidal natural isomorphism. This means $\varphi = \{\varphi_c : c \rightarrow c^{\vee\vee}\}$ is a collection of natural isomorphisms such that for all $a, b \in \mathcal{C}$, the following diagram commutes:

$$\begin{array}{ccc} a \otimes b & \xrightarrow{\varphi_a \otimes \varphi_b} & (a \otimes b)^{\vee\vee} \\ \downarrow \varphi_{a \otimes b} & & \downarrow (\nu_{b^\vee, a^\vee})^\vee \\ a^{\vee\vee} \otimes b^{\vee\vee} & \xrightarrow{\nu_{a^\vee, b^\vee}} & (b^\vee \otimes a^\vee)^\vee \end{array} \quad (4)$$

A *pivotal category* is a rigid monoidal category equipped with a pivotal structure.

Two pivotal structures (\vee_1, φ^1) and (\vee_2, φ^2) on \mathcal{C} are *equivalent* if for every $c \in \mathcal{C}$, the following diagram commutes:

$$\begin{array}{ccc} c & \xrightarrow{\varphi_c^1} & c_1^{\vee\vee} \\ & \searrow \varphi_c^2 & \uparrow \\ & & c_2^{\vee\vee} \end{array} \quad \int_{c_2^{\vee\vee}}^{\text{coev}_{c_2}} \int_{c_1^{\vee\vee}}^{\text{ev}_{c_1}} \quad (5)$$

⁷We use the notation of [DSPS13]; \mathcal{C}^{mop} denotes the category obtained from \mathcal{C} by reversing arrows and reversing the order of tensor product. In other words, $\vee : \mathcal{C} \rightarrow \mathcal{C}$ is contravariant and anti-monoidal.

Remark 2.5. If \mathcal{C} has a pivotal structure, then the equivalence classes of pivotal structures on \mathcal{C} form a torsor for the group $\text{Aut}_{\otimes}(\text{id}_{\mathcal{C}})$ of monoidal natural automorphisms of the identity functor of \mathcal{C} [EGNO15, Ex. 4.7.16].

Definition 2.6. Given a pivotal category (\mathcal{C}, φ) , we define the left and right trace on $\mathcal{C}(c \rightarrow c)$ for each $c \in \mathcal{C}$ by

$$\text{tr}_L^{\varphi}(f) := c^{\vee} \left(\begin{array}{c} \boxed{f} \\ \boxed{\varphi_c^{-1}} \end{array} \right)_{c^{\vee\vee}} \quad \text{tr}_R^{\varphi}(f) := \left(\begin{array}{c} \boxed{\varphi_c} \\ \boxed{f} \end{array} \right)_{c^{\vee}} . \quad (6)$$

2.2 Semisimple pivotal categories For this section, (\mathcal{C}, φ) is a semisimple pivotal multi-tensor category. This means $\mathcal{C}(1_{\mathcal{C}} \rightarrow 1_{\mathcal{C}})$ is a finite dimensional complex semisimple algebra, and is thus isomorphic to \mathbb{C}^r for some $r \in \mathbb{N}$. The next lemma is well known to experts; we include a proof for convenience and completeness.

Lemma 2.7. *The traces tr_L^{φ} and tr_R^{φ} are nondegenerate, i.e., for every nonzero $f \in \mathcal{C}(a \rightarrow b)$, there is a $g \in \mathcal{C}(b \rightarrow a)$ such that $\text{tr}_L(g \circ f) \neq 0$, and similarly for tr_R .*

Proof. Suppose $f \in \mathcal{C}(a \rightarrow b)$ is nonzero. Then there is a simple $c \in \mathcal{C}$, a monomorphism $g \in \mathcal{C}(c \rightarrow a)$, and an epimorphism $h \in \mathcal{C}(b \rightarrow c)$ such that $h \circ f \circ g \neq 0$. Then

$$0 \neq e := \text{ev}_c \circ [\text{id}_{c^{\vee}} \otimes (h \circ f \circ g \circ \varphi_c^{-1})] \in \mathcal{C}(c^{\vee} \otimes c^{\vee\vee} \rightarrow 1_{\mathcal{C}}).$$

Since we also know $0 \neq \text{coev}_{c^{\vee}} \in \mathcal{C}(1_{\mathcal{C}} \rightarrow c^{\vee} \otimes c^{\vee\vee})$, by [HPT16a, Lem. A.5],

$$\text{tr}_L((g \circ h) \circ f) = \text{tr}_L(h \circ f \circ g) = e \circ \text{coev}_{c^{\vee}} \neq 0.$$

Hence tr_L is nondegenerate. The proof that tr_R is nondegenerate is similar and left to the reader. \square

Definition 2.8. Let $1_{\mathcal{C}} = \bigoplus_{i=1}^r 1_i$ be a decomposition into simples, and for $1 \leq i \leq r$, let $p_i \in \mathcal{C}(1_{\mathcal{C}} \rightarrow 1_{\mathcal{C}})$ be the minimal idempotent corresponding to 1_i . For $c \in \mathcal{C}$ and $f \in \mathcal{C}(c \rightarrow c)$, we define the $M_r(\mathbb{C})$ -valued traces Tr_L^{φ} and Tr_R^{φ} by the formulas

$$\begin{aligned} (\text{Tr}_L^{\varphi}(f))_{i,j} \text{id}_{1_j} &= \text{tr}_L^{\varphi}(p_i \otimes f \otimes p_j) = c^{\vee} \left(\begin{array}{c} \boxed{f} \\ \boxed{\varphi_c^{-1}} \end{array} \right)_{c^{\vee\vee}} \\ (\text{Tr}_R^{\varphi}(f))_{i,j} \text{id}_{1_i} &= \text{tr}_R^{\varphi}(p_i \otimes f \otimes p_j) = \left(\begin{array}{c} \boxed{\varphi_c} \\ \boxed{f} \end{array} \right)_{c^{\vee}} \end{aligned} \quad (7)$$

Notice that $\mathrm{Tr}_L^\varphi(f)^T = \mathrm{Tr}_R^\varphi(f^\vee)$ and $\mathrm{Tr}_L^\varphi(f^\vee) = \mathrm{Tr}_R^\varphi(c)^T$ for all $c \in \mathcal{C}$ and $f \in \mathcal{C}(c \rightarrow c)$. Moreover, $\mathrm{Tr}_L^\varphi, \mathrm{Tr}_R^\varphi : \mathcal{C}(c \rightarrow c) \rightarrow M_r(\mathbb{C})$ are tracial; for all $f \in \mathcal{C}(c \rightarrow d)$ and $g \in \mathcal{C}(d \rightarrow c)$, we have $\mathrm{Tr}_L^\varphi(g \circ f) = \mathrm{Tr}_L^\varphi(f \circ g)$, and similarly for Tr_R^φ .

We call (\mathcal{C}, φ) *spherical* if for every $c \in \mathcal{C}$ and $f \in \mathrm{End}_{\mathcal{C}}(c)$, $\mathrm{Tr}_L^\varphi(f) = \mathrm{Tr}_R^\varphi(f)$.

For each $c \in \mathcal{C}$, we define $\mathrm{Dim}_L^\varphi(c), \mathrm{Dim}_R^\varphi(c) \in M_r(\mathbb{C})$ by

$$\mathrm{Dim}_L^\varphi(c) := \mathrm{Tr}_L^\varphi(\mathrm{id}_c) \quad \mathrm{Dim}_R^\varphi(c) := \mathrm{Tr}_R^\varphi(c). \quad (8)$$

Notice that $\mathrm{Dim}_L^\varphi(c)^T = \mathrm{Dim}_R^\varphi(c^\vee)$ and $\mathrm{Dim}_L^\varphi(c^\vee) = \mathrm{Dim}_R^\varphi(c)^T$ for all $c \in \mathcal{C}$. Moreover, $\mathrm{Dim}_L^\varphi, \mathrm{Dim}_R^\varphi : K_0(\mathcal{C}) \rightarrow M_r(\mathbb{C})$ are ring homomorphisms. For each simple $c \in \mathcal{C}$, the matrices $\mathrm{Dim}_L^\varphi(c)$ and $\mathrm{Dim}_R^\varphi(c)$ have exactly one non-zero entry, which we denote $\mathrm{dim}_L^\varphi(c)$ and $\mathrm{dim}_R^\varphi(c)$ respectively.

Corollary 2.9 ([EGNO15, Prop. 4.8.4]). *For all simple $c \in \mathcal{C}$, $\mathrm{dim}_L^\varphi(c) \neq 0 \neq \mathrm{dim}_R^\varphi(c)$.*

Definition 2.10. A pivotal structure (\vee, φ) on a semisimple multitensor category \mathcal{C} is called *pseudounitary* if $\mathrm{dim}_L^\varphi(c) > 0$ and $\mathrm{dim}_R^\varphi(c) > 0$ for all simple $c \in \mathrm{Irr}(\mathcal{C})$. This definition is equivalent to [EGNO15, Def. 9.4.4] in the context of fusion categories by uniqueness of the Frobenius-Perron dimensions.

Remark 2.11. Suppose \mathcal{C} is a semisimple multitensor category which has a pseudounitary pivotal structure. Then similar to Remark 2.5, the equivalence classes of pseudounitary pivotal structures on \mathcal{C} forms a torsor for the subgroup $\mathrm{Aut}_{\otimes}^+(\mathrm{id}_{\mathcal{C}})$ of $\mathrm{Aut}_{\otimes}(\mathrm{id}_{\mathcal{C}})$ of *positive* monoidal natural automorphisms of the identity dagger tensor functor, which consists of those monoidal natural isomorphisms $\zeta : \mathrm{id}_{\mathcal{C}} \Rightarrow \mathrm{id}_{\mathcal{C}}$ such that for every simple $c \in \mathcal{C}$, $\zeta_c : c \rightarrow c$ is a *strictly positive* multiple of id_c .

Lemma 2.12. *For two pivotal structures (\vee_1, φ^1) and (\vee_2, φ^2) , the following are equivalent:*

- (1) (\vee_1, φ^1) and (\vee_2, φ^2) are equivalent.
- (2) For all $c \in \mathcal{C}$ and $f \in \mathcal{C}(c \rightarrow c)$, $\mathrm{tr}_L^1(f) = \mathrm{tr}_L^2(f)$.
- (3) For all $c \in \mathcal{C}$ and $f \in \mathcal{C}(c \rightarrow c)$, $\mathrm{tr}_R^1(f) = \mathrm{tr}_R^2(f)$.
- (4) For all simple $c \in \mathcal{C}$, $\mathrm{dim}_L^1(c) = \mathrm{dim}_L^2(c)$.
- (5) For all simple $c \in \mathcal{C}$, $\mathrm{dim}_R^1(c) = \mathrm{dim}_R^2(c)$.

Proof.

(1) \Rightarrow (2): This is straightforward.

(2) \Leftrightarrow (3): Note that $\mathrm{tr}_L^1(f) = \mathrm{tr}_R^1(f^\vee)$ and $\mathrm{tr}_L^2(f) = \mathrm{tr}_R^2(f^\vee)$ for all $f \in \mathcal{C}(c \rightarrow c)$.

(2) \Rightarrow (4): Take $f = \mathrm{id}_c$.

(4) \Leftrightarrow (5): Note that $\mathrm{dim}_L^1(c) = \mathrm{dim}_R^1(c_1^\vee)$ and $\mathrm{dim}_L^2(c) = \mathrm{dim}_R^2(c_2^\vee)$ for all simple $c \in \mathcal{C}$.

(4) \Rightarrow (1): By monoidality of φ^1, φ^2 , and the canonical intertwining morphism in (5), (\vee_1, φ^1) and (\vee_2, φ^2) are equivalent if and only if for all simple $c \in \mathcal{C}$,

$$(\varphi_c^2)^{-1} \circ \left(\text{redrawing of } \varphi_c^1 \right) \circ \varphi_c^1 = \mathrm{id}_c. \quad (9)$$

Now the left hand side of (9) is a scalar multiple of id_c . By Corollary 2.9, we may determine this scalar by applying tr_L^1 to both sides as $\mathrm{dim}_L^i(c) \neq 0$ for $i = 1, 2$. It is straightforward to check that tr_L^1 applied to the left hand side is equal to $\mathrm{dim}_L^2(c)$, which is equal to $\mathrm{dim}_L^1(c)$ by assumption. Hence (9) holds. \square

2.3 Pivotal functors

Definition 2.13. Given a strong monoidal functor between pivotal categories $(F, \mu) : (\mathcal{C}, \varphi^{\mathcal{C}}) \rightarrow (\mathcal{D}, \varphi^{\mathcal{D}})$, for each c in \mathcal{C} , we have a canonical natural isomorphism $\delta_c : F(c^\vee) \rightarrow F(c)^\vee$ given by

$$\delta_c := \begin{array}{c} \boxed{F(\text{ev}_c)} \\ \downarrow \text{ } F(c^\vee \otimes c) \quad \downarrow \text{ } F(c)^\vee \\ \boxed{\mu_{F(c^\vee), F(c)}} \\ \downarrow \text{ } F(c^\vee) \quad \downarrow \text{ } F(c) \end{array} = (F(\text{ev}_c) \otimes \text{id}_{F(c)^\vee}) \circ (\mu_{F(c^\vee), F(c)} \otimes \text{id}_{F(c)^\vee}) \circ (\text{id}_{F(c^\vee)} \otimes \text{coev}_{F(c)}). \quad (10)$$

We call (F, μ) *pivotal* if for all $c \in \mathcal{C}$,

$$\begin{array}{c} F(c^\vee)^\vee \\ \downarrow \text{ } \delta_c^\vee \\ F(c)^{\vee\vee} \\ \downarrow \text{ } \varphi_{F(c)}^{\mathcal{D}} \\ F(c) \end{array} = \begin{array}{c} F(c^\vee)^\vee \\ \downarrow \text{ } \delta_{c^\vee} \\ F(c^{\vee\vee}) \\ \downarrow \text{ } \varphi_{F(c)}^{\mathcal{C}} \\ F(c) \end{array}. \quad (11)$$

Lemma 2.14. Suppose $(\mathcal{C}, \varphi^{\mathcal{C}})$ and $(\mathcal{D}, \varphi^{\mathcal{D}})$ are pivotal categories and $(F, \mu) : \mathcal{C} \rightarrow \mathcal{D}$ is a pivotal strong monoidal functor. Then (F, μ) preserves the left and right pivotal traces, i.e., for all $f \in \mathcal{C}(c \rightarrow c)$,

$$F \left(\begin{array}{c} c \\ \downarrow \text{ } f \\ c \\ \downarrow \text{ } \varphi_c^{-1} \\ c^{\vee\vee} \end{array} \right) = F(\text{tr}_L^{\varphi^{\mathcal{C}}}(f)) = \text{tr}_L^{\varphi^{\mathcal{D}}}(F(f)) = F(c)^\vee \begin{array}{c} F(c) \\ \downarrow \text{ } F(f) \\ F(c) \\ \downarrow \text{ } \varphi_{F(c)}^{-1} \\ F(c)^{\vee\vee} \end{array}, \quad (12)$$

and similarly for the right pivotal traces.

Proof. Notice that for $f \in \mathcal{C}(c \rightarrow c)$, we always have

$$F(\text{tr}_L^{\mathcal{C}}(f)) = \begin{array}{c} \boxed{F(\text{ev}_c)} \\ \downarrow \text{ } F(c^\vee \otimes c) \\ \boxed{\mu} \\ \downarrow \text{ } F(c) \\ \boxed{F(f)} \\ \downarrow \text{ } F(c) \\ \boxed{F(\varphi_c^{-1})} \\ \downarrow \text{ } F(c^{\vee\vee}) \\ \boxed{\mu^{-1}} \\ \downarrow \text{ } F(c^\vee \otimes c^{\vee\vee}) \\ \boxed{F(\text{coev}_{c^\vee})} \end{array} = \begin{array}{c} F(c)^\vee \quad \downarrow \text{ } \delta_c \quad \downarrow \text{ } F(f) \\ \downarrow \text{ } F(c) \\ \boxed{F(\varphi_c^{-1})} \\ \downarrow \text{ } F(c^{\vee\vee}) \\ \boxed{\delta_{c^\vee}^{-1}} \\ \downarrow \text{ } F(c^\vee)^\vee \end{array} = \begin{array}{c} F(c)^\vee \quad \downarrow \text{ } F(f) \\ \downarrow \text{ } F(c) \\ \boxed{F(\varphi_c^{-1})} \\ \downarrow \text{ } F(c^{\vee\vee}) \\ \boxed{\delta_{c^\vee}^{-1}} \\ \downarrow \text{ } F(c^\vee)^\vee \\ \boxed{\delta_c^\vee} \\ \downarrow \text{ } F(c)^{\vee\vee} \end{array} \quad (13)$$

If (F, μ) is pivotal, then the right hand sides of (12) and (13) above are equal. The proof for the right pivotal trace is analogous. \square

The converse of Lemma 2.14 is true under some additional assumptions.

Lemma 2.15. *If $(\mathcal{C}, \varphi^{\mathcal{C}}), (\mathcal{D}, \varphi^{\mathcal{D}})$ are pivotal semisimple multitensor categories and $(F, \mu) : \mathcal{C} \rightarrow \mathcal{D}$ is a full strong monoidal functor which preserves the left or right pivotal traces, then (F, μ) is fully faithful and pivotal.*

Proof. We assume (F, μ) preserves the left pivotal traces, and the proof for the right pivotal traces is analogous. First, suppose $f \in \mathcal{C}(c \rightarrow d)$. By nondegeneracy of $\text{tr}_L^{\varphi^{\mathcal{C}}}$, there is a $g \in \mathcal{C}(d \rightarrow c)$ such that $\text{tr}_L^{\varphi^{\mathcal{C}}}(g \circ f) \neq 0$. Then $\text{tr}_L^{\varphi^{\mathcal{D}}}(F(g) \circ F(f)) \neq 0$, so $F(f) \neq 0$ and F is fully faithful. Notice this immediately implies F takes simples to simples, and non-isomorphic simples in \mathcal{C} remain non-isomorphic in \mathcal{D} . Now to show (F, μ) is pivotal, by monoidality, it suffices to prove (11) when $c \in \mathcal{C}$ is simple. By the above argument, $F(c)$ is then simple, and every morphism in $\mathcal{D}(F(c) \rightarrow F(c))$ is a scalar multiple of $\text{id}_{F(c)}$. Since the right hand side of (12) is equal to the right hand side of (13) by hypothesis, by nondegeneracy of the trace from Lemma 2.7, we must have $\varphi_{F(c)}^{-1} = F(\varphi_c^{-1}) \circ \delta_{c^\vee}^{-1} \circ \delta_c^\vee$, and thus (11) holds. \square

3. Unitary dual functors

We begin this section with background on dagger structures and \mathbb{C}^* categories in §3.1. Next, we give the correct notion of unitary dual functor and unitary pivotal structure from [Sel11, §7.3]. Similar to the situation for tensor categories, in §3.3, we classify pivotal structures on a multitensor category via homomorphisms out of the universal grading groupoid \mathcal{U} . In §3.4, for a unitary multitensor category, we construct a canonical unitary dual functor from each groupoid homomorphism $\pi \in \text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$, and we show these exhaust the possible unitary equivalence classes of unitary dual functors. We then describe the canonical bi-involutive structure associated to a unitary dual functor in §3.5.

3.1 Dagger structures and unitary multitensor categories

Definition 3.1. Given a \mathbb{C} -linear category \mathcal{C} , a *dagger structure* is a collection of anti-linear maps $\dagger : \mathcal{C}(c \rightarrow d) \rightarrow \mathcal{C}(d \rightarrow c)$ for all $c, d \in \mathcal{C}$ such that $(f \circ g)^\dagger = g^\dagger \circ f^\dagger$ for composable f and g , and $f^{\dagger\dagger} = f$ for all f . A morphism $f : \mathcal{C}(a \rightarrow b)$ is called *unitary* if $f^\dagger = f^{-1}$.

A *dagger (multi)tensor category* is a (multi)tensor category equipped with a dagger structure so that $(f \otimes g)^\dagger = f^\dagger \otimes g^\dagger$ for all morphisms f, g , and all associator and unitors are unitary natural isomorphisms.

Definition 3.2. A functor between dagger categories $F : \mathcal{M} \rightarrow \mathcal{N}$ is called a *dagger functor* if $F(f^\dagger) = F(f)^\dagger$ for all morphisms f in \mathcal{M} . Given finitely semisimple dagger categories \mathcal{M} and \mathcal{N} , we define $\text{Fun}^\dagger(\mathcal{M} \rightarrow \mathcal{N})$ to be the dagger category of dagger functors from $\mathcal{C} \rightarrow \mathcal{D}$ with dagger structure defined as follows. Given a natural transformation of dagger functors $\eta : F \Rightarrow G$, it is straightforward to show that $(\eta^\dagger)_m := (\eta_m)^\dagger$ for $m \in \mathcal{M}$ gives a well-defined natural transformation $\eta^\dagger : G \Rightarrow F$.⁸ One now calculates that $\eta \mapsto \eta^\dagger$ defines a dagger structure on $\text{Fun}^\dagger(\mathcal{M} \rightarrow \mathcal{N})$. It is important to note that a natural transformation $\eta : F \Rightarrow G$ is unitary if and only if $\eta_m \in \mathcal{N}(F(m) \rightarrow G(m))$ is unitary for all $m \in \mathcal{M}$.

A *dagger equivalence* of dagger categories \mathcal{M} and \mathcal{N} consists of dagger functors $F : \mathcal{M} \rightarrow \mathcal{N}$ and $G : \mathcal{N} \rightarrow \mathcal{M}$ together with unitary natural isomorphisms $\text{id} \Rightarrow F \circ G$ and $\text{id} \Rightarrow G \circ F$. A tensor functor between dagger tensor categories $(F, \mu) : \mathcal{C} \rightarrow \mathcal{D}$ is called a *dagger tensor functor* if F is a dagger functor and $\mu_{c,d}$ is unitary for all $c, d \in \mathcal{C}$. Given a finitely semisimple dagger

⁸In the \mathbb{C}^* setting, one only considers *bounded* natural transformations, i.e., those for which $\sup_{m \in \mathcal{M}} \|\eta_m\| < \infty$.

category \mathcal{M} , $\text{End}^\dagger(\mathcal{M}) := \text{Fun}^\dagger(\mathcal{M}, \mathcal{M})$ is easily seen to be a strict semisimple dagger multitensor category.

Remark 3.3. The *principle of equivalence* [nLa18] in category theory roughly states that a properly defined structure should be invariant under the proper notion of equivalence. The proper notion of equivalence between two objects in a dagger category is that they are *unitary isomorphic*, and the proper notion of equivalence between dagger functors between dagger categories is that they are *unitarily naturally isomorphic*. For example, if $F, G : \mathcal{C} \rightarrow \mathcal{D}$ are functors between dagger categories with F a dagger functor, and $\eta : F \Rightarrow G$ is a natural isomorphism, G need not be a dagger functor unless η is unitary.

With this in mind, a dagger category *cannot* be considered as a category with some extra categorical structure. For example, if \mathcal{D} is a dagger category and the underlying category of \mathcal{D} is equivalent to the category \mathcal{C} , there is generally no way to endow \mathcal{C} with a dagger structure which promotes the equivalence to a dagger equivalence. We refer the reader to the helpful discussion between Shulman and Selinger available at [nLa18] for further details.

Remark 3.4. The forgetful functor on a finitely semisimple dagger (multitensor) category which forgets the dagger structure is a fully faithful (tensor) functor. Thus the category Hilb_{fd} of finite dimensional Hilbert spaces is equivalent to the category Vec_{fd} of finite dimensional vector spaces as a linear (tensor) category. Also, for a finitely semisimple dagger category \mathcal{M} , $\text{End}^\dagger(\mathcal{M}) \cong \text{End}(\mathcal{M})$ as linear multitensor categories.

Definition 3.5. A dagger category which admits orthogonal direct sums is called a C^* category if every endomorphism algebra is a C^* -algebra (see [GLR85, HP17]). Notice that a finitely semisimple dagger category is C^* if and only if it is dagger equivalent to Hilb^n for some $n \in \mathbb{N}$.

Definition 3.6. A *tensor C^* category* is a linear monoidal dagger category which admits orthogonal direct sums and is idempotent complete, and whose underlying dagger category is C^* . A *unitary (multi)tensor category* is a semisimple (multi)tensor C^* category.⁹ As before, the prefix *multi-* is used if and only if $1_{\mathcal{C}}$ is not simple.

A *unitary (multi)fusion category* is a finitely semisimple unitary (multi)tensor category. For $r > 1$, an $r \times r$ *unitary multifusion category* is an indecomposable unitary multifusion category such that $\dim(\mathcal{C}(1_{\mathcal{C}} \rightarrow 1_{\mathcal{C}})) = r$, so we can orthogonally decompose $1_{\mathcal{C}}$ into simples as $\bigoplus_{i=1}^r 1_i$.

Warning 3.7. While natural from a *non-unitary* viewpoint, pseudounitary pivotal structures are *unnatural* for unitary multitensor categories. The problem arises from the fact that while a dual functor is unique up to unique natural isomorphism as in (3), this unique natural isomorphism need not be unitary! Hence one may only discuss the compatibility of a *fixed* dual functor $\vee : \mathcal{C} \rightarrow \mathcal{C}^{\text{mop}}$ with \dagger . However, compatibility of \dagger with \vee need not imply compatibility of \dagger with an equivalent dual functor $\vee' : \mathcal{C} \rightarrow \mathcal{C}^{\text{mop}}$. We provide Lemma 3.12 below which gives a sufficient condition to transport the compatibility. In §3.4 below, we give a *manifestly unitary* approach, and we reconcile the latter with the former.

Lemma 3.8. *Suppose \mathcal{C} is a unitary multitensor category and φ is a pivotal structure. The following are equivalent.*

⁹By a generalization of [LR97, Lem. 3.9] (see also the second paragraph on p. 9 therein), a tensor C^* category is rigid if and only if it is semisimple. Hence the adjective ‘tensor’ in ‘tensor C^* category’ does *not* include ‘rigid’ nor having simple unit object, in conflict with the definition of ‘tensor category’ following [EGNO15].

(1) (\mathcal{C}, φ) is pseudounitary.

(2) For all $a, b \in \mathcal{C}$ and all $f \in \mathcal{C}(a \rightarrow b)$ with $f \neq 0$, $\mathrm{tr}_L(f^\dagger \circ f) > 0$ and $\mathrm{tr}_R(f^\dagger \circ f) > 0$.¹⁰

Proof. That (2) \Rightarrow (1) is trivial. Suppose (\mathcal{C}, φ) is pseudounitary. We show that for all $a, b \in \mathcal{C}$ and $f \in \mathcal{C}(a \rightarrow b)$ with $f \neq 0$, $\mathrm{tr}_L^\varphi(f^\dagger \circ f) > 0$. The proof that $\mathrm{tr}_R^\varphi(f \circ f^\dagger) > 0$ is similar.

Step 1: Suppose $c \in \mathcal{C}$ is simple and $f \in \mathcal{C}(c \rightarrow c)$ with $f \neq 0$. Then $f = \lambda \mathrm{id}_c$ for some $\lambda \in \mathbb{C}^\times$, so $f^\dagger \circ f = |\lambda|^2 \mathrm{id}_c$, and $\mathrm{tr}_L^\varphi(f^\dagger \circ f) = |\lambda|^2 \dim_L^\varphi(c) > 0$.

Step 2: Suppose $a, b \in \mathcal{C}$ are respectively orthogonal direct sums of m, n objects isomorphic to the simple object $c \in \mathcal{C}$ and $f \in \mathcal{C}(a \rightarrow b)$ with $f \neq 0$. Pick m isometries $v_1, \dots, v_m \in \mathcal{C}(c \rightarrow a)$ so that $\sum_{i=1}^m v_i \circ v_i^\dagger = \mathrm{id}_a$. Note that $f \neq 0$ if and only if $v_i^\dagger \circ f^\dagger \circ f \circ v_i \in \mathcal{C}(c \rightarrow c)$ is nonzero for some $i = 1, \dots, m$. Thus by Step 1,

$$\mathrm{tr}_L^\varphi(f^\dagger \circ f) = \sum_{i=1}^m \mathrm{tr}_L^\varphi(v_i \circ v_i^\dagger \circ f^\dagger \circ f) = \sum_{i=1}^m \mathrm{tr}_L^\varphi(v_i^\dagger \circ f^\dagger \circ f \circ v_i) > 0.$$

Step 3: For arbitrary $a, b \in \mathcal{C}$ and $f \in \mathcal{C}(a \rightarrow b)$ nonzero, decompose a and b into orthogonal direct sums of isotypic components and apply Step 2. \square

3.2 Unitary dual functors

For this section, \mathcal{C} is a unitary multitensor category. The following proposition is [Sel11, Lem. 7.5], which can be viewed as a generalization of [Vic11, Lem. 2.16] in the non-strict unitary multitensor category setting.

Proposition 3.9. Fix a dual functor $\vee : \mathcal{C} \rightarrow \mathcal{C}^{\mathrm{mop}}$ with its canonical tensorator ν from (2). The following are equivalent.

- (1) \vee is a dagger tensor functor, i.e., for all $a, b \in \mathcal{C}$ and $f \in \mathcal{C}(a \rightarrow b)$, $\nu_{a,b}$ is unitary and $f^{\vee\dagger} = f^{\dagger\vee}$.
- (2) Defining $\varphi_c := (\mathrm{coev}_c^\dagger \otimes \mathrm{id}_{c^{\vee\vee}}) \circ (\mathrm{id}_c \otimes \mathrm{coev}_{c^\vee})$ gives a pivotal structure $\varphi : \mathrm{id} \Rightarrow \vee \circ \vee$.

Proof. First, note that φ is natural if and only if

$$\varphi_b \circ f = f^{\vee\vee} \circ \varphi_a \quad \forall f \in \mathcal{C}(a \rightarrow b)$$

if and only if

$$\mathrm{coev}_b^\dagger \circ (f \otimes \mathrm{id}_{b^\vee}) = \mathrm{coev}_a^\dagger \circ (\mathrm{id}_a \otimes f^\vee) \quad \forall f \in \mathcal{C}(a \rightarrow b)$$

if and only if $f^{\dagger\vee} = f^{\vee\dagger}$ for all $f \in \mathcal{C}(a \rightarrow b)$. Second, note that φ is monoidal if and only if

$$\mathrm{coev}_a^\dagger \circ (\mathrm{id}_a \otimes \mathrm{coev}_b^\dagger \otimes \mathrm{id}_{a^\vee}) = \mathrm{coev}_{a \otimes b}^\dagger \circ (\mathrm{id}_{a \otimes b} \otimes \nu_{b,a}) \quad \forall a, b \in \mathcal{C}$$

if and only if

$$\mathrm{id}_{b^\vee \otimes a^\vee} = (\mathrm{id}_{b^\vee \otimes a^\vee} \otimes \mathrm{coev}_{a \otimes b}^\dagger) \circ (\mathrm{id}_{b^\vee} \otimes \mathrm{ev}_a^\dagger \otimes \mathrm{id}_b \otimes \mathrm{id}_{(a \otimes b)^\vee}) \circ (\mathrm{ev}_b^\dagger \otimes \mathrm{id}_{(a \otimes b)^\vee}) \circ \nu_{b,a} \quad \forall a, b \in \mathcal{C}$$

if and only if $\nu_{a,b}$ is unitary for all $a, b \in \mathcal{C}$. \square

Corollary 3.10. If either of the equivalent conditions of Proposition 3.9 hold, then for all $c \in \mathcal{C}$,

$$(\mathrm{coev}_c^\dagger \otimes \mathrm{id}_{c^{\vee\vee}}) \circ (\mathrm{id}_c \otimes \mathrm{coev}_{c^\vee}) =: \varphi_c = (\mathrm{id}_{c^{\vee\vee}} \otimes \mathrm{ev}_c) \circ (\mathrm{ev}_{c^\vee}^\dagger \otimes \mathrm{id}_c), \quad (14)$$

which is equivalent to φ_c being unitary for all $c \in \mathcal{C}$.

¹⁰Note that if $f^{\vee\dagger} = f^{\dagger\vee}$, then $\mathrm{tr}_L(f^\dagger \circ f) > 0$ if and only if $\mathrm{tr}_R(f^\dagger \circ f) > 0$.

Proof. By (1) of Proposition 3.9, we have $\text{coev}_c^{\dagger\vee} = \text{coev}_c^{\vee\dagger} = \text{ev}_{c^\vee}^\dagger$, which is equivalent to (14). Now notice that φ_c^\dagger is the inverse of the expression on the right hand side of (14), so (14) holds if and only if φ_c is unitary. \square

Definition 3.11. A dual functor $\vee : \mathcal{C} \rightarrow \mathcal{C}^{\text{mop}}$ is called a *unitary dual functor* if any of the equivalent conditions of Proposition 3.9 hold. Two unitary dual functors are called *unitarily equivalent* if the canonical monoidal natural isomorphism from (3) is unitary.

In line with the principle of equivalence for dagger categories discussed in Remark 3.3, unitary equivalence between a unitary dual functor and an arbitrary dual functor transports unitarity, as we will see right below in Lemma 3.12. Of course, two unitary dual functors need not be unitarily equivalent, as can be seen from the construction in §3.4 together with Lemma 3.15 below.

Lemma 3.12. *Suppose $\nu_1, \nu_2 : \mathcal{C} \rightarrow \mathcal{C}^{\text{mop}}$ are two dual functors such that \vee_1 is unitary. If for all $c \in \mathcal{C}$, the canonical isomorphism $\zeta_c \in \mathcal{C}(c^{\vee_2} \rightarrow c^{\vee_1})$ from (3) is unitary, then \vee_2 is unitary.*

Proof. Suppose that the canonical isomorphism in (3) is always unitary. Then for all $f \in \mathcal{C}(a \rightarrow b)$,

$$\begin{aligned} f^{\vee_2\dagger} &= (\text{id}_{b^{\vee_2}} \otimes (\text{coev}_a^2)^\dagger) \circ (\text{id}_{b^{\vee_2}} \otimes f^\dagger \otimes \text{id}_{a^{\vee_2}}) \circ ((\text{ev}_b^2)^\dagger \otimes \text{id}_{a^{\vee_2}}) \\ &= (\text{id}_{b^{\vee_2}} \otimes (\text{coev}_b^1)^\dagger \otimes (\text{coev}_a^1)^\dagger \otimes (\text{coev}_a^2)^\dagger) \circ (\text{id}_{b^{\vee_2} \otimes b \otimes b^{\vee_1}} \otimes f^\dagger \otimes \text{id}_{a^{\vee_1} \otimes a \otimes a^{\vee_2}}) \\ &\quad \circ ((\text{ev}_b^2)^\dagger \otimes (\text{ev}_b^1)^\dagger \otimes (\text{ev}_a^1)^\dagger \otimes \text{id}_{a^{\vee_2}}) \\ &= (\text{ev}_a^2 \otimes \text{id}_{a^{\vee_1}}) \circ (\text{id}_{a^{\vee_2}} \otimes \text{coev}_a^1) \circ f^{\dagger\vee_1} \circ (\text{ev}_b^2 \otimes \text{id}_{b^{\vee_1}}) \circ (\text{id}_{b^{\vee_2}} \otimes \text{coev}_b^1) \\ &= f^{\dagger\vee_2}. \end{aligned}$$

Moreover, for all $a, b \in \mathcal{C}$, we have

$$\nu_{a,b}^2 = \zeta_{b \otimes a}^{-1} \circ \nu_{a,b}^1 \circ (\zeta_a \otimes \zeta_b) \in \mathcal{C}(a^{\vee_2} \otimes b^{\vee_2} \rightarrow (b \otimes a)^{\vee_2}),$$

which is necessarily unitary as it is a composite of unitaries. Hence \vee_2 is unitary. \square

Definition 3.13. We call a pivotal structure (\vee, φ) *unitary* if \vee is a unitary dual functor and φ is as in (14). Two unitary pivotal structures are *unitarily equivalent* if they are equivalent and the canonical monoidal natural isomorphism from (5) is unitary.

Remark 3.14. For a unitary dual functor \vee , the left and right pivotal traces have alternate formulas that show they are manifestly positive linear operators $\mathcal{C}(c \rightarrow c) \rightarrow \mathcal{C}(1_{\mathcal{C}} \rightarrow 1_{\mathcal{C}})$:

$$\begin{aligned} \text{tr}_L^\varphi(f) &\stackrel{(6)}{=} \text{ev}_c \circ (\text{id}_{c^\vee} \otimes f) \circ (\text{id}_{c^\vee} \otimes \varphi_c^{-1}) \circ \text{coev}_{c^\vee} \stackrel{\text{Cor. 3.10}}{=} \text{ev}_c \circ (\text{id}_{c^\vee} \otimes f) \circ \text{ev}_c^\dagger \\ \text{tr}_R^\varphi(f) &\stackrel{(6)}{=} \text{ev}_c \circ (\varphi_c \otimes \text{id}_{c^\vee}) \circ (f \otimes \text{id}_{c^\vee}) \circ \text{coev}_{c^\vee} \stackrel{\text{Cor. 3.10}}{=} \text{coev}_c^\dagger \circ (f \otimes \text{id}_{c^\vee}) \circ \text{coev}_c. \end{aligned}$$

Hence every unitary pivotal structure is pseudounitary. We will use these alternate formulas in §3.4 below.

Lemma 3.15. *Fix two unitary dual functors \vee_1, \vee_2 , and let φ^1, φ^2 be the respective induced unitary pivotal structures. We have \vee_1 and \vee_2 are unitarily equivalent if and only if φ^1 and φ^2 are unitarily equivalent.*

Proof. Recall that for $c \in \mathcal{C}$, $\zeta_c \in \mathcal{C}(c^{\vee 2} \rightarrow c^{\vee 1})$ is the unique natural isomorphism from (3). Observe that \vee_1 and \vee_2 are unitarily equivalent if and only if

$$\zeta_c^{-1} = (\mathrm{id}_{c^{\vee 2}} \otimes \mathrm{coev}_1^\dagger) \circ (\mathrm{ev}_2^\dagger \otimes \mathrm{id}_{c^{\vee 1}}) = \zeta_c^\dagger \quad \forall c \in \mathcal{C}$$

if and only if

$$\mathcal{J} = \varphi_c^2 \circ (\varphi_c^1)^{-1} = (\mathrm{coev}_1^\dagger \otimes \mathrm{id}_{c^{\vee 1 \vee 1}}) \circ (\mathrm{id}_c \otimes \mathrm{coev}_{c^{\vee 1}}) \circ (\mathrm{ev}_{c^{\vee 2}} \otimes \mathrm{id}_c) \circ (\mathrm{id}_{c^{\vee 2 \vee 2}} \otimes \mathrm{ev}_2^\dagger) \quad \forall c \in \mathcal{C}$$

if and only if φ^1 and φ^2 are unitarily equivalent. \square

3.3 The universal grading groupoid We now adapt [EGNO15, §4.14] to the (unitary) multitensor category setting. For this section, \mathcal{C} is a multitensor category which is not necessarily unitary. In the unitary setting, one should add the terms in parentheses, and one may ignore them in the algebraic setting.

Recall that a *groupoid* \mathcal{G} is a category where all morphisms are invertible. We will identify \mathcal{G} with its set of morphisms, and we can recover the objects as the *idempotents*, i.e., those morphisms $e \in \mathcal{G}$ such that $e \circ e = e$.

Definition 3.16. A *grading* of \mathcal{C} by a groupoid \mathcal{G} is a decomposition

$$\mathcal{C} = \bigoplus_{g \in \mathcal{G}} \mathcal{C}_g$$

where each $\mathcal{C}_g \subset \mathcal{C}$ is a semisimple (\mathbb{C}^*) subcategory such that if g, h are composable, the tensor product maps $\mathcal{C}_g \times \mathcal{C}_h$ to \mathcal{C}_{gh} . When g, h are not composable, the tensor product on $\mathcal{C}_g \times \mathcal{C}_h$ is the zero bi-functor. For every idempotent in $e \in \mathcal{G}$, \mathcal{C}_e is a (unitary) multitensor subcategory of \mathcal{C} . A grading by \mathcal{G} is called *faithful* if $\mathcal{C}_g \neq 0$ for all $g \in \mathcal{G}$. Gradings are in bijection with gradings of the Grothendieck ring $K_0(\mathcal{C})$ as a based ring, where the basis corresponds to the isomorphism classes of simple objects of \mathcal{C} .

Given any two faithful gradings, there is a common faithful refinement, so there exists a universal grading of \mathcal{C} . We call the groupoid associated to the universal grading of \mathcal{C} the *universal grading groupoid*, denoted \mathcal{U} .

Remark 3.17. If \mathcal{C} is faithfully graded by \mathcal{G} , then since \mathcal{U} is a refinement of \mathcal{G} , we get a canonical surjective groupoid homomorphism $\mathcal{U} \rightarrow \mathcal{G}$ given by mapping a $u \in \mathcal{U}$ to the $g \in \mathcal{G}$ such that $\mathcal{C}_u \subset \mathcal{C}_g$.

Recall that when \mathcal{C} is a tensor category, the universal grading group can be obtained as the quotient of the free group $\mathbb{F}[\mathrm{Irr}(\mathcal{C})]$ by relations of the form $c = ab$ whenever c is isomorphic to a summand of $a \otimes b$. For a multitensor category with $\dim(\mathrm{End}(1_{\mathcal{C}})) = r$, the universal grading groupoid can be obtained as follows. First, choose a decomposition $1_{\mathcal{C}} = \bigoplus_{i=1}^r 1_i$ into simples, and let $\mathcal{C}_{ij} := 1_i \otimes \mathcal{C} \otimes 1_j$ be the ij -th component. Choose a set of representatives of simple objects $\mathrm{Irr}(\mathcal{C}_{ij})$ of \mathcal{C}_{ij} . We define \mathcal{F} to be the free groupoid with r objects $1, \dots, r$ and morphisms generated by the simples $\mathrm{Irr}(\mathcal{C}_{ij}) \subset \mathrm{Hom}(i \rightarrow j)$ for all $1 \leq i, j \leq r$. We impose the relations $c = ab$ whenever simples a and b are composable in \mathcal{F} and the simple c is isomorphic to a summand of $a \otimes b$.

Remark 3.18. Observe that when \mathcal{C} is an $r \times r$ multitensor category (indecomposable with $\dim(\text{End}_{\mathcal{C}}(1_{\mathcal{C}})) = r$), then the universal grading groupoid \mathcal{U} is always (non-canonically) isomorphic to $\mathcal{M}_r \times G$ for some group G . Moreover, G is no larger than the universal grading group of \mathcal{C}_{ii} for any $1 \leq i \leq r$.


Example 3.19. The matrix category $\text{Mat}_n(\text{Vec})$ has universal grading groupoid \mathcal{M}_n which has n objects and a unique isomorphism between any two objects. We can identify the morphism set of this groupoid with the standard system of matrix units $\{E_{ij}\}$ for $M_n(\mathbb{C})$. Given a tensor category \mathcal{C} and $n \in \mathbb{N}$, $\text{Mat}_n(\mathcal{C})$ has universal grading groupoid isomorphic to $\mathcal{M}_n \times \mathcal{U}_{\mathcal{C}}$ where $\mathcal{U}_{\mathcal{C}}$ is the universal grading group of \mathcal{C} .

Example 3.20. Let G be a finite group. The multifusion category

$$\begin{pmatrix} \text{Vec}(G) & \text{Vec} \\ \text{Vec} & \text{Rep}(G) \end{pmatrix}$$

has universal grading groupoid \mathcal{M}_2 .

Example 3.21. Consider the TLJ category \mathcal{A}_{17} and let $a = \mathbf{1} + \mathbf{9} + \mathbf{17} \in \mathcal{A}_{17}$ be the algebra object whose category of modules is $\mathcal{E}_7 = \text{Mod}_{\mathcal{A}_{17}}(a)$. The fusion rules for the generators m, m' of the dual category \mathcal{B} of $a - a$ bimodules are given by the following fusion diagram taken from [HPT16b, Ex. 3.17] (see also [Got10, §5.3.5]):



The multifusion category

$$\begin{pmatrix} \mathcal{A}_{17} & \mathcal{E}_7 \\ \mathcal{E}_7^{\text{op}} & \mathcal{B} \end{pmatrix}$$

has universal grading groupoid isomorphic to $\mathcal{M}_2 \times \mathbb{Z}/2\mathbb{Z}$. This follows from Remark 3.18 together with the fact that the universal grading groups of \mathcal{A}_{17} and \mathcal{B} are both $\mathbb{Z}/2\mathbb{Z}$, and that the module category \mathcal{E}_7 has a compatible $\mathbb{Z}/2\mathbb{Z}$ -grading.

Example 3.22. The Haagerup subfactor gives 2 exotic fusion categories which have trivial universal grading groups [AH99]. There is one other fusion category in this Morita equivalence class [GS12], and the three fusion categories can be combined into one 3×3 multifusion category as in [GMP⁺18, Prop. 3.10]. This 3×3 Haagerup multifusion category has universal grading groupoid \mathcal{M}_3 .

Similarly, there is a 4×4 Extended Haagerup multifusion category with universal grading groupoid \mathcal{M}_4 [GMP⁺18].

Example 3.23 (Alternating part). Given a tensor category \mathcal{C} and an $X \in \mathcal{C}$ which Cauchy tensor generates, the *alternating part* is the 2×2 multitensor subcategory of $\text{Mat}_2(\mathcal{C})$ generated by $X \in \mathcal{C}_{12}$ and $\overline{X} \in \mathcal{C}_{21}$. By Remark 3.18, the universal grading groupoid of the alternating part

is no larger than $\mathcal{M}_2 \times G_{\text{ad}}$ where G_{ad} is the universal grading group of the adjoint subcategory of \mathcal{C} .

For an explicit example, the *standard invariant* multitensor subcategory of the $A_{-\infty, \infty}$ subfactor $N \subset M$ [Jon83, Pop94]

$$\begin{pmatrix} \text{Bim}(N) & \text{Bim}(N, M) \\ \text{Bim}(M, N) & \text{Bim}(M) \end{pmatrix} = \langle L^2 M \rangle \subset \text{Bim}(N \oplus M)$$

is the alternating part of $\text{Mat}_2(\text{Hilb}_{\text{fd}}(\mathbb{Z}))$, which has universal grading groupoid isomorphic to $\mathcal{M}_2 \times \mathbb{Z}$.

Notation 3.24. For $c \in \mathcal{C}$, we say c is *homogeneous* if c lies in one component subcategory $\mathcal{C}_g \subset \mathcal{C}$ for some $g \in \mathcal{U}$. For such c , we define $\text{gr}(c) := g$. For an arbitrary $c \in \mathcal{C}$, we write $c = \bigoplus_{g \in \mathcal{U}} c_g$ for the canonical (orthogonal) decomposition of c into homogeneous subobjects. For an $f \in \mathcal{C}(a \rightarrow b)$, we write $f_g \in \mathcal{C}(a_g \rightarrow b_g)$ for the g -graded component of f .

For a groupoid \mathcal{G} and an abelian group A (whose group law is still denoted multiplicatively), we denote by $\text{Hom}(\mathcal{G} \rightarrow A)$ the set of functors from \mathcal{G} to A where the latter is viewed as a groupoid with exactly one object. Note that $\text{Hom}(\mathcal{G} \rightarrow A)$ is a group under pointwise multiplication and pointwise inversion.

Recall from Remarks 2.5 and 2.11 that $\text{Aut}_{\otimes}(\text{id}_{\mathcal{C}})$ is the group of monoidal natural automorphisms of the identity tensor functor, and $\text{Aut}_{\otimes}^+(\text{id}_{\mathcal{C}})$ is the subgroup of positive monoidal natural isomorphisms of the identity dagger tensor functor.

Lemma 3.25. *There is a canonical isomorphism $\text{Aut}_{\otimes}(\text{id}_{\mathcal{C}}) \cong \text{Hom}(\mathcal{U} \rightarrow \mathbb{C}^{\times})$ which takes the subgroup $\text{Aut}_{\otimes}^+(\text{id}_{\mathcal{C}})$ onto the subgroup $\text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$.*

Proof. Given $\zeta \in \text{Aut}_{\otimes}(\text{id}_{\mathcal{C}})$, we get a grading of \mathcal{C} by \mathbb{C}^{\times} by assigning to each simple $c \in \mathcal{C}$ the number corresponding to $\zeta_c \in \mathcal{C}(c \rightarrow c) = \mathbb{C} \text{id}_c$. This gives us a homomorphism $f_{\zeta} : \mathcal{U} \rightarrow \mathbb{C}^{\times}$ by universality of \mathcal{U} . One now checks the map $\zeta \mapsto f_{\zeta}$ is an isomorphism. Finally, $\zeta \in \text{Aut}_{\otimes}^+(\text{id}_{\mathcal{C}})$ if and only if $\zeta_c \in \mathbb{R}_{>0} \text{id}_c$ for all simple $c \in \mathcal{C}$ if and only if $\text{im}(f_{\zeta}) \subset \mathbb{R}_{>0}$. \square

For the convenience of the reader, we provide the lemma below which, in the presence of a pseudounitary pivotal structure (\vee, φ) , provides an explicit bijection between the torsor of pseudounitary pivotal structures on \mathcal{C} and $\text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$ obtained by combining Remark 2.11 with Lemma 3.25.

Lemma 3.26. *Suppose (\vee, φ) is a pivotal structure on a semisimple multitensor category \mathcal{C} . Defining for a simple $c \in \mathcal{C}$*

$$\pi(\text{gr}(c)) := \frac{\dim_L^{\varphi}(c)}{\dim_R^{\varphi}(c)} \tag{15}$$

gives a well defined a homomorphism $\pi : \mathcal{U} \rightarrow \mathbb{C}^{\times}$. If (\vee, φ) is pseudounitary, then $\text{im}(\pi) \subset \mathbb{R}_{>0}$.

Proof. First, note that for a simple $c \in \mathcal{C}$, $\dim_L^{\varphi}(c) \neq 0 \neq \dim_R^{\varphi}(c)$ by Corollary 2.9. Next, if $c, d \in \mathcal{C}$ are simple such that $c \otimes d \neq 0$,

$$\text{Dim}_L^{\varphi}(c) \text{Dim}_L^{\varphi}(d) = \text{Dim}_L^{\varphi}(c \otimes d) \implies \dim_L^{\varphi}(c \otimes d) = \dim_L^{\varphi}(c) \dim_L^{\varphi}(d), \tag{16}$$

and similarly for the right dimension. If moreover $\text{gr}(c) = \text{gr}(d)$, then $e := \text{gr}(c \otimes d^\vee)$ is an idempotent in \mathcal{U} (an identity morphism). Since (\vee, φ) restricted to \mathcal{C}_e gives a spherical tensor category, we have

$$\dim_L^\varphi(c \otimes d^\vee) = \dim_R^\varphi(c \otimes d^\vee) \iff \frac{\dim_L^\varphi(c)}{\dim_R^\varphi(c)} = \frac{\dim_L^\varphi(d)}{\dim_R^\varphi(d)},$$

and π is well-defined. Now (16) immediately implies that π is a homomorphism. The last claim is obvious. \square

3.4 Balanced duals In this section, \mathcal{C} is a unitary multitensor category. The following definition was suggested by André Henriques.

Definition 3.27. Let $\pi : \mathcal{U} \rightarrow \mathbb{R}_{>0}$ be a groupoid homomorphism. Denote by Ψ the linear functional in $\mathcal{C}(1 \rightarrow 1) \rightarrow \mathbb{C}$ which sends every minimal projection to $1_{\mathbb{C}}$. A π -balanced dual of $c \in \mathcal{C}$ is a triple $(c^\vee, \text{ev}_c^\pi, \text{coev}_c^\pi)$ such that the morphisms $\text{ev}_c^\pi, \text{coev}_c^\pi$ satisfy the zig-zag axioms and the π -balancing condition: for all $f \in \mathcal{C}(c \rightarrow c)$ and $g \in \mathcal{U}$

$$\Psi \left(\text{ev}_c^\pi \circ (\text{id}_{c^\vee} \otimes f g) \circ (\text{ev}_c^\pi)^\dagger \right) = \pi(g) \cdot \Psi \left((\text{coev}_c^\pi)^\dagger \circ (f g \otimes \text{id}_{c^\vee}) \circ \text{coev}_c^\pi \right). \quad (17)$$

A unitary dual functor \vee_π is called π -balanced if (17) holds for all $c \in \mathcal{C}$, $f \in \mathcal{C}(c \rightarrow c)$, and $g \in \mathcal{U}$.

If $\pi(g) = 1$ for all $g \in \mathcal{U}$, we omit π from the notation; we simply say $(c^\vee, \text{ev}_c, \text{coev}_c)$ is a balanced dual which satisfies the zig-zag axioms and the balancing condition. A dual functor \vee is balanced if (17) holds with $\pi(g) = 1$ for all $c \in \mathcal{C}$, $f \in \mathcal{C}(c \rightarrow c)$, and $g \in \mathcal{U}$.

Our next task is to construct a π -balanced unitary dual functor for every $\pi \in \text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$. To do so, we will use the following facts from [Yam04].

Fact 3.28 ([Yam04, Lem. 3.6]). *Suppose $(c^\vee, \text{ev}_c, \text{coev}_c)$ is an arbitrary dual of $c \in \mathcal{C}$. The positive map $\mathcal{C}(c \rightarrow c) \rightarrow \mathcal{C}(1 \rightarrow 1)$ given by $f \mapsto \text{ev}_c \circ (\text{id}_{c^\vee} \otimes f) \circ \text{ev}_c^\dagger$ is faithful: for any $g \in \mathcal{C}(c \rightarrow d)$,*

$$\text{ev}_c \circ (\text{id}_{c^\vee} \otimes (g^\dagger \circ g)) \circ \text{ev}_c^\dagger = 0 \iff (\text{id}_{c^\vee} \otimes g) \circ \text{ev}_c^\dagger = 0 \iff g = 0.$$

Similarly, $f \mapsto \text{coev}_c^\dagger \circ (f \otimes \text{id}_{c^\vee}) \circ \text{coev}_c$ is faithful.¹¹

Fact 3.29 ([Yam04, Lem. 3.7.ii], [BKLR15, Prop. 2.6]). *For $a, b \in \mathcal{C}$ with arbitrary choices of duals $(a^\vee, \text{ev}_a, \text{coev}_a)$, $(b^\vee, \text{ev}_b, \text{coev}_b)$ respectively, the following are equivalent:*

- (1) For all $f \in \mathcal{C}(a \rightarrow b)$, $f^{\vee\dagger} = f^{\dagger\vee}$.
- (2) For all $g \in \mathcal{C}(b \rightarrow a)$, $g^{\vee\dagger} = g^{\dagger\vee}$.
- (3) For all $a, b \in \mathcal{C}$, $f \in \mathcal{C}(a \rightarrow b)$, and $g \in \mathcal{C}(b \rightarrow a)$,

$$\text{ev}_a \circ (\text{id}_{a^\vee} \otimes (g \circ f)) \circ \text{ev}_a^\dagger = \text{ev}_b \circ (\text{id}_{b^\vee} \otimes (f \circ g)) \circ \text{ev}_b^\dagger. \quad (18)$$

- (4) For all $a, b \in \mathcal{C}$, $f \in \mathcal{C}(a \rightarrow b)$, and $g \in \mathcal{C}(b \rightarrow a)$,

$$\text{coev}_a^\dagger \circ ((g \circ f) \otimes \text{id}_{a^\vee}) \circ \text{coev}_a = \text{coev}_b^\dagger \circ ((f \circ g) \otimes \text{id}_{b^\vee}) \circ \text{coev}_b. \quad (19)$$

¹¹More is true: given a morphism $\varepsilon \in \mathcal{C}(c^\vee \otimes c \rightarrow 1_{\mathcal{C}})$, the map $f \mapsto \varepsilon \circ (\text{id}_{c^\vee} \otimes f) \circ \varepsilon^\dagger$ is faithful if and only if there is a morphism $\eta \in \mathcal{C}(1_{\mathcal{C}} \rightarrow c \otimes c^\vee)$ such that $(c^\vee, \varepsilon, \eta)$ is a dual of c . This is proven in [Yam04, Lem. 3.6] where the condition that $1_{\mathcal{C}}$ is simple is never used. A similar statement holds swapping ε and η .

Proof.

(1) \Rightarrow (2): Suppose (1) holds and $g \in \mathcal{C}(b \rightarrow a)$. Then $g^\dagger \in \mathcal{C}(a \rightarrow b)$, so

$$g^{\dagger\vee} = (g^\dagger)^{\vee\dagger\dagger} = (g^\dagger)^{\dagger\vee\dagger} = g^{\vee\dagger}.$$

(2) \Rightarrow (1): Analogous to (1) \Rightarrow (2).

(2) \Leftrightarrow (3): This is similar to the proof of [Yam04, Lem. 3.7.ii], which does not require that $1_{\mathcal{C}}$ is simple or that $a = b$. (Indeed, [BKLR15, Prop. 2.6] does not assume $a = b$.) In more detail, observe that for $f \in \mathcal{C}(a \rightarrow b)$ and $g \in \mathcal{C}(b \rightarrow a)$,

$$\text{ev}_a \circ (\text{id}_{a^\vee} \otimes (g \circ f)) \circ \text{ev}_a^\dagger = \text{ev}_b \circ (g^\vee \otimes f) \circ \text{ev}_a^\dagger.$$

By faithfulness from Fact 3.28, the above is equal to the right hand side of (18) if and only if

$$(\text{coev}_b^\dagger \otimes \text{id}_a) \circ (\text{id}_b \otimes g^\vee \otimes \text{id}_a) \circ (\text{id}_b \otimes \text{ev}_a^\dagger) = g \quad \iff \quad g^{\vee\dagger} = g^{\dagger\vee}.$$

(1) \Leftrightarrow (4): This is similar to (2) \Leftrightarrow (3) and left to the reader. \square

Proposition 3.30. *Fix a groupoid homomorphism $\pi : \mathcal{U} \rightarrow \mathbb{R}_{>0}$. For each $c \in \mathcal{C}$, there exists a unique π -balanced dual $(c^{\vee\pi}, \text{ev}_c^\pi, \text{coev}_c^\pi)$ up to unique unitary isomorphism.*

Proof. We adapt the proof from [Yam04, Lem. 3.9] and [NT13, Prop. 2.2.15].

Step 1: Suppose $c \in \mathcal{C}$ is simple, and let $(c^\vee, \text{ev}_c, \text{coev}_c)$ be a dual of c . Then (17) is satisfied if and only if $\Psi(\text{ev}_c \circ \text{ev}_c^\dagger) = \pi(\text{gr}(c))\Psi(\text{coev}_c^\dagger \circ \text{coev}_c)$. Since $\pi(\text{gr}(c)) > 0$, we can scale ev_c and coev_c by inverse scalars to achieve this. Moreover, this choice of scalar is unique up to a phase. Hence the choice of ev_c^π and coev_c^π satisfying (17) is unique up to a unique phase in $U(1)$.

Step 2: Suppose $c \in \mathcal{C}$ is an orthogonal direct sum of n objects isomorphic to the simple object $a \in \mathcal{C}$. Let $(a^{\vee\pi}, \text{ev}_a^\pi, \text{coev}_a^\pi)$ be the unique π -balanced dual of a from Step 1. Suppose $c^{\vee\pi} \in \mathcal{C}$ can be equipped with an evaluation and coevaluation which make it a dual for c . Pick n isometries $v_1, \dots, v_n \in \mathcal{C}(a \rightarrow c)$ and $w_1, \dots, w_n \in \mathcal{C}(a^{\vee\pi} \rightarrow c^{\vee\pi})$ with orthogonal ranges so that $\sum_{i=1}^n v_i \circ v_i^\dagger = \text{id}_c$ and $\sum_{i=1}^n w_i \circ w_i^\dagger = \text{id}_{c^{\vee\pi}}$. Define

$$\text{ev}_c^\pi := \sum_{i=1}^n \text{ev}_a^\pi \circ (w_i^\dagger \otimes v_i^\dagger) \quad \text{and} \quad \text{coev}_c^\pi := \sum_{i=1}^n (v_i \otimes w_i) \circ \text{coev}_a^\pi.$$

It is clear that ev_c^π and coev_c^π satisfy the zig-zag axioms. Moreover, for $v_k \circ v_\ell^\dagger$, we calculate that

$$\begin{aligned} & \text{ev}_c^\pi \circ (\text{id}_{c^{\vee\pi}} \otimes (v_k \circ v_\ell^\dagger)) \circ (\text{ev}_c^\pi)^\dagger \\ &= \sum_{i,j=1}^n \text{ev}_a^\pi \circ (w_i^\dagger \otimes v_i^\dagger) \circ (\text{id}_{c^{\vee\pi}} \otimes (v_k \circ v_\ell^\dagger)) \circ (w_j \otimes v_j) \circ (\text{ev}_a^\pi)^\dagger \\ &= \sum_{i,j=1}^n \text{ev}_a^\pi \circ (\text{id}_{c^{\vee\pi}} \otimes (v_i^\dagger \circ v_k \circ v_\ell^\dagger \circ v_j)) \circ (\text{ev}_a^\pi)^\dagger \\ &= \delta_{k=\ell} (\text{ev}_a^\pi \circ (\text{ev}_a^\pi)^\dagger) \end{aligned} \tag{20}$$

and similarly,

$$(\text{coev}_c^\pi)^\dagger \circ ((v_k \circ v_\ell^\dagger) \otimes \text{id}_{c^{\vee\pi}}) \circ \text{coev}_c^\pi = \delta_{k=\ell} ((\text{coev}_a^\pi)^\dagger \circ \text{coev}_a^\pi). \tag{21}$$

Hence (17) is satisfied since it was true for $a \in \mathcal{C}$ by Step 1.

Now suppose in addition to $(c^{\vee\pi}, \text{ev}_c^\pi, \text{coev}_c^\pi)$, we have a second dual $(c^\vee, \text{ev}_c, \text{coev}_c)$ which is π -balanced. Pick n isometries $x_1, \dots, x_n \in \mathcal{C}(a \rightarrow c)$ and $y_1, \dots, y_n \in \mathcal{C}(a^{\vee\pi} \rightarrow c^\vee)$ with orthogonal ranges so that $\sum_{i=1}^n x_i \circ x_i^\dagger = \text{id}_c$ and $\sum_{i=1}^n y_i \circ y_i^\dagger = \text{id}_{c^\vee}$. Since $(c^\vee, \text{ev}_c, \text{coev}_c)$ is π -balanced, so is $(a^\vee, \text{ev}_c \circ (y_i \otimes x_i), (x_i^\dagger \otimes y_i^\dagger) \circ \text{coev}_c)$, so by Step 1, there is a unique phase $\phi_i \in U(1)$ such that $\text{ev}_a^\pi = \phi_i \cdot \text{ev}_c \circ (y_i \otimes x_i)$ and $\text{coev}_a^\pi = \overline{\phi_i} \cdot (x_i^\dagger \otimes y_i^\dagger) \circ \text{coev}_c$. We replace y_i with $\phi_i \cdot y_i$ for each i so that $\text{ev}_a^\pi = \text{ev}_c \circ (y_i \otimes x_i)$ and $\text{coev}_a^\pi = (x_i^\dagger \otimes y_i^\dagger) \circ \text{coev}_c$.

Now $\mathcal{C}(c \rightarrow c) \cong M_n(\mathbb{C})$ acts on the Hilbert space $\mathcal{C}(a \rightarrow c) \cong \mathbb{C}^n$ with the isometry inner product determined by the formula $\langle f, g \rangle_{\text{Isom}} \text{id}_a = g^\dagger \circ f$ by post-composition. Since $\{x_i\}_{i=1}^n$ and $\{y_i\}_{i=1}^n$ are orthonormal bases for this inner product, there is a unitary $u \in \mathcal{C}(c \rightarrow c)$ such that $ux_i = y_i$ for $i = 1, \dots, n$. Define the scalars $u_{ij} := x_i^\dagger \circ u \circ x_j \in \mathcal{C}(a \rightarrow a) \cong \mathbb{C}$, and observe that $(u_{ij})_{i,j=1}^n \in M_n(\mathbb{C})$ is unitary. Define $U := \sum_{i,j} u_{ij}(w_j \circ y_i^\dagger) \in \mathcal{C}(c^\vee \rightarrow c^{\vee\pi})$, which is necessarily unitary by construction. Then we have

$$v_j = u \circ x_j = \sum_i u_{ij} x_i \quad \text{and} \quad U \circ y_i = \sum_j u_{ij} w_j,$$

which implies

$$\sum_i x_i \otimes_{\mathbb{C}} (U \circ y_i) = \sum_{i,j} u_{i,j} (x_i \otimes_{\mathbb{C}} w_j) = \sum_j v_j \otimes_{\mathbb{C}} w_j \in \mathcal{C}(a \rightarrow c) \otimes_{\mathbb{C}} \mathcal{C}(a^\vee \rightarrow c^{\vee\pi}).$$

Now applying the map $f \otimes_{\mathbb{C}} g \mapsto (f \otimes g) \circ \text{coev}_a^\pi \in \mathcal{C}(1_c \rightarrow c \otimes c^{\vee\pi})$, we see

$$(\text{id}_c \otimes U) \circ \text{coev}_c = \left(\sum_i x_i \otimes (U \circ y_i) \right) \circ \text{coev}_a^\pi = \left(\sum_j v_j \otimes w_j \right) \circ \text{coev}_a^\pi = \text{coev}_c^\pi.$$

By Remark 2.3, we have that the unique isomorphism $c^\vee \rightarrow c^{\vee\pi}$ is equal to U and is necessarily unitary.

Step 3: Suppose $c \in \mathcal{C}$ is arbitrary. Decompose c into an orthogonal direct sum of isotypic components and apply Step 2. \square

Lemma 3.31. *Fix a groupoid homomorphism $\pi : \mathcal{U} \rightarrow \mathbb{R}_{>0}$. Consider π -balanced duals $(c^{\vee\pi}, \text{ev}_c^\pi, \text{coev}_c^\pi)$ and $(d^{\vee\pi}, \text{ev}_d^\pi, \text{coev}_d^\pi)$ of c and d respectively. For $c \otimes d \in \mathcal{C}$, the dual*

$$(d^{\vee\pi} \otimes c^{\vee\pi}, \text{ev}_d^\pi \circ (\text{id}_{d^{\vee\pi}} \otimes \text{ev}_c^\pi \otimes \text{id}_d), (\text{id}_c \otimes \text{coev}_d^\pi \otimes \text{id}_{c^{\vee\pi}}) \circ \text{coev}_c^\pi)$$

is π -balanced.

Proof. We prove the lemma in the case c, d are homogeneous following [Yam04, Lem. 3.11.i], and we leave the rest of the details to the reader. In this case, for $f \in \mathcal{C}(c \otimes d \rightarrow c \otimes d)$, f is homogeneous, and we define

$$\begin{aligned} E_c^L(f) &:= (\text{ev}_c^\pi \otimes \text{id}_d) \circ (\text{id}_{c^{\vee\pi}} \otimes f) \circ ((\text{ev}_c^\pi)^\dagger \otimes \text{id}_d) \in \mathcal{C}(d \rightarrow d) \\ E_d^R(f) &:= (\text{id}_c \otimes (\text{coev}_d^\pi)^\dagger) \circ (f \otimes \text{id}_{d^{\vee\pi}}) \circ (\text{id}_c \otimes \text{coev}_d^\pi) \in \mathcal{C}(d \rightarrow d), \end{aligned}$$

which are also homogeneous morphisms. We then calculate

$$\begin{aligned}
& \Psi \left((\text{ev}_d^\pi \circ (\text{id}_{d^{\vee\pi}} \otimes \text{ev}_c^\pi \otimes \text{id}_d)) \circ (\text{id}_{d^{\vee\pi} \otimes c^{\vee\pi}} \otimes f) \circ (\text{ev}_d^\pi \circ (\text{id}_{d^{\vee\pi}} \otimes \text{ev}_c^\pi \otimes \text{id}_d))^\dagger \right) \\
&= \Psi \left(\text{ev}_d^\pi \circ (\text{id}_{d^{\vee\pi}} \otimes E_c^L(f)) \circ (\text{ev}_d^\pi)^\dagger \right) \\
&= \pi(\text{gr}(d)) \cdot \Psi \left((\text{coev}_d^\pi)^\dagger \circ (E_c^L(f) \otimes \text{id}_{d^{\vee\pi}}) \circ \text{coev}_d^\pi \right) \\
&= \pi(\text{gr}(d)) \cdot \Psi \left(\text{ev}_c^\pi \circ (\text{id}_{c^{\vee\pi}} \otimes E_d^R(f)) \circ (\text{ev}_c^\pi)^\dagger \right) \\
&= \pi(\text{gr}(c))\pi(\text{gr}(d)) \cdot \Psi \left((\text{coev}_c^\pi)^\dagger \circ (\text{id}_{c^{\vee\pi}} \otimes E_d^R(f)) \circ \text{coev}_c^\pi \right) \\
&= \pi(\text{gr}(c \otimes d)) \cdot \\
& \quad \Psi \left(((\text{id}_c \otimes \text{coev}_d^\pi \otimes \text{id}_{c^{\vee\pi}}) \circ \text{coev}_c^\pi)^\dagger \circ (f \otimes \text{id}_{d^{\vee\pi} \otimes c^{\vee\pi}}) \circ ((\text{id}_c \otimes \text{coev}_d^\pi \otimes \text{id}_{c^{\vee\pi}}) \circ \text{coev}_c^\pi) \right).
\end{aligned}$$

Thus the dual $(d^{\vee\pi} \otimes c^{\vee\pi}, \text{ev}_d^\pi \circ (\text{id}_{d^{\vee\pi}} \otimes \text{ev}_c^\pi \otimes \text{id}_d), (\text{id}_c \otimes \text{coev}_d^\pi \otimes \text{id}_{c^{\vee\pi}}) \circ \text{coev}_c^\pi)$ is π -balanced. \square

Corollary 3.32. *For every $\pi \in \text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$, there exists a unique π -balanced unitary dual functor \vee_π up to unique unitary monoidal natural isomorphism.*

Proof. By Proposition 3.30, for every $c \in \mathcal{C}$, there is a unique π -balanced dual $(c^{\vee\pi}, \text{ev}_c^\pi, \text{coev}_c^\pi)$ up to unique unitary isomorphism. By Lemma 3.31 and Proposition 3.30, the canonical tensorator ν^π from (2) is necessarily unitary. It remains to prove \vee_π is a dagger functor. As in the proof of [Yam04, Lem. 3.9], By (20) and (21), equations (18) and (19) hold, i.e., the positive linear maps

$$\begin{aligned}
f &\mapsto \text{ev}_c^\pi \circ (\text{id}_{c^{\vee\pi}} \otimes f) \circ (\text{ev}_c^\pi)^\dagger \in \mathcal{C}(1 \rightarrow 1) \\
f &\mapsto (\text{coev}_c^\pi)^\dagger \circ (f \otimes \text{id}_{c^{\vee\pi}}) \circ \text{coev}_c^\pi \in \mathcal{C}(1 \rightarrow 1)
\end{aligned}$$

are tracial (and faithful by Fact 3.28). Hence by Fact 3.29, \vee_π is a dagger functor. \square

Theorem (Theorem A). *Let \mathcal{C} be a unitary multitensor category. There are canonical bijections between:*

- (1) Pseudounitary pivotal structures up to monoidal natural isomorphism.
- (2) Unitary dual functors up to unitary monoidal natural isomorphism.
- (3) $\text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$.

Proof.

(2) \Leftrightarrow (3): Suppose \vee is a unitary dual functor, and let φ be the canonical associated unitary pivotal structure, which is pseudounitary by Remark 3.14. Thus (15) from Lemma 3.26 gives us a function $\vee \mapsto \pi_\vee$ from unitary equivalence classes of unitary dual functors to $\text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$, which is injective by Lemmas 2.12 and 3.15. Surjectivity now follows immediately from Corollary 3.32, since it is easy to calculate that $\pi_{\vee_\pi} = \pi$ by (17).

(1) \Leftrightarrow (3): Since a pseudounitary pivotal structure exists on \mathcal{C} , this follows immediately by combining Remark 2.11 and Lemma 3.25. \square

Remark 3.33. Suppose \mathcal{C} is faithfully graded by the groupoid \mathcal{G} . Then for any $\pi \in \text{Hom}(\mathcal{G} \rightarrow \mathbb{R}_{>0})$, we get a unique lift $\tilde{\pi} \in \text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$ using the canonical canonical groupoid surjection $\mathcal{U} \twoheadrightarrow \mathcal{G}$ from Remark 3.17. Then the unique $\tilde{\pi}$ -balanced unitary dual functor is π -balanced: for all simple $c \in \mathcal{C}$ with $\text{gr}(c) = g \in \mathcal{G}$ and all $f \in \mathcal{C}(c \rightarrow c)$,

$$\Psi(\text{ev}_c^{\tilde{\pi}} \circ (\text{id}_{c^\vee} \otimes f) \circ (\text{ev}_c^{\tilde{\pi}})^\dagger) = \pi(g) \cdot \Psi((\text{coev}_c^{\tilde{\pi}})^\dagger \circ (f \otimes \text{id}_{c^\vee}) \circ \text{coev}_c^{\tilde{\pi}}).$$

Choosing \mathcal{G} to be trivial or $\pi \in \text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$ trivial yields the following corollary.

Corollary 3.34 ([Yam04, Thm. 4.7] and [BDH14, §4]). *A unitary multitensor category has a unique unitary spherical structure corresponding to $\pi = 1$ such that for all $f \in \mathcal{C}(c \rightarrow c)$ and $g \in \mathcal{U}$,*

$$\Psi \left(\text{coev}_c^\dagger \circ (f_g \otimes \text{id}_{c^\vee}) \circ \text{coev}_c \right) = \Psi \left(\text{ev}_c \circ (\text{id}_{c^\vee} \otimes f_g) \circ \text{ev}_c^\dagger \right). \quad (22)$$

Remark 3.35. Starting with a balanced unitary dual functor \vee , we can *rescale* \vee by a $\pi \in \text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$ to obtain a π -balanced unitary dual functor $\vee_\pi : \mathcal{C} \rightarrow \mathcal{C}^{\text{mop}}$ as follows. For a homogeneous $c \in \mathcal{C}$ with $\text{gr}(c) = g \in \mathcal{U}$, we define

$$\text{ev}_c^\pi := \pi(g)^{1/4} \text{ev}_c \quad \text{coev}_c^\pi := \pi(g)^{-1/4} \text{coev}_c. \quad (23)$$

It is immediate that these renormalized maps satisfy the zig-zag axioms, and moreover, we see $(c^\vee, \text{ev}_c^\pi, \text{coev}_c^\pi)$ is π -balanced:

$$\begin{aligned} \text{ev}_c^\pi \circ (\text{id}_{c^\vee} \otimes f) \circ (\text{ev}_c^\pi)^\dagger &= \pi(g)^{1/2} \cdot \left(\text{ev}_c \circ (\text{id}_{c^\vee} \otimes f) \circ \text{ev}_c^\dagger \right) \\ &= \pi(g)^{1/2} \cdot \left(\text{coev}_c^\dagger \circ (f \otimes \text{id}_{c^\vee}) \circ \text{coev}_c \right) \\ &= \pi(g) \cdot \left((\text{coev}_c^\pi)^\dagger \circ (f \otimes \text{id}_{c^\vee}) \circ \text{coev}_c^\pi \right). \end{aligned}$$

However, notice that we have left ν and φ *unchanged!* Indeed, if c, d are homogeneous with $\text{gr}(c) = g$ and $\text{gr}(d) = h$, then $c \otimes d$ is homogeneous with $\text{gr}(c \otimes d) = gh$, and

$$\begin{aligned} \nu_{c,d}^\pi &= (\text{ev}_d^\pi \otimes \text{id}_{(c \otimes d)^\vee}) \circ (\text{id}_{d^\vee} \otimes \text{ev}_c^\pi \otimes \text{id}_d \otimes \text{id}_{(c \otimes d)^\vee}) \circ (\text{id}_{d^\vee \otimes c^\vee} \otimes \text{coev}_{c \otimes d}^\pi) \\ &= \pi(g)^{1/4} \pi(h)^{1/4} \pi(gh)^{-1/4} \cdot \\ &\quad \left((\text{ev}_d \otimes \text{id}_{(c \otimes d)^\vee}) \circ (\text{id}_{d^\vee} \otimes \text{ev}_c \otimes \text{id}_d \otimes \text{id}_{(c \otimes d)^\vee}) \circ (\text{id}_{d^\vee \otimes c^\vee} \otimes \text{coev}_{c \otimes d}) \right) \\ &= \nu_{c,d}. \end{aligned}$$

Similarly, $\text{gr}(c^\vee) = g^{-1}$, and

$$\begin{aligned} \varphi_c^\pi &= ((\text{coev}_c^\pi)^\dagger \otimes \text{id}_{c^{\vee\vee}}) \circ (\text{id}_c \otimes \text{coev}_{c^\vee}^\pi) \\ &= \pi(g)^{-1/4} \pi(g^{-1})^{-1/4} \cdot \left((\text{coev}_c^\dagger \otimes \text{id}_{c^{\vee\vee}}) \circ (\text{id}_c \otimes \text{coev}_{c^\vee}) \right) = \varphi_c. \end{aligned}$$

Since ν, φ are completely determined on homogeneous objects by naturality, we see that $\nu^\pi = \nu$ and $\varphi^\pi = \varphi$.

Conversely, starting with \vee_π a π -balanced unitary dual functor, we can obtain a balanced unitary dual functor by rescaling evaluations and coevaluations on homogeneous objects $c \in \mathcal{C}$ with $\text{gr}(c) = g \in \mathcal{U}$ by

$$\text{ev}_c := \pi(g)^{-1/4} \text{ev}_c^\pi \quad \text{coev}_c := \pi(g)^{1/4} \text{coev}_c^\pi. \quad (24)$$

As before, ν and φ remained unchanged by this scaling.

3.5 Bi-involutive structures The notion of unitary dual functor is stronger than the similar notion of *bi-involutive structure* from [HP17, §2.1].

Definition 3.36. An *involutive structure* [Egg11] on a multitensor category \mathcal{C} consists of a conjugate-linear tensor functor $(\bar{\cdot}, \nu) : \mathcal{C} \rightarrow \mathcal{C}^{\text{mp}}$ ¹² together with a monoidal natural isomorphism $\varphi : \text{id}_{\mathcal{C}} \rightarrow \bar{\bar{\cdot}}$ ¹³. When \mathcal{C} is unitary, we call $(\bar{\cdot}, \nu, \varphi)$ a *bi-involutive structure* [HP17] if $\bar{\cdot}$ is a dagger functor and ν, φ are unitary.

Example 3.37. Complex conjugation gives a bi-involutive structure on the tensor \mathbf{C}^* category Hilb of separable Hilbert spaces.

Example 3.38. Similar to the previous example, complex conjugation gives a bi-involutive structure on $\text{Bim}(M)$, the tensor \mathbf{C}^* category of separable $M - M$ bimodules where M is any von Neumann algebra with the Connes fusion tensor product.

By Proposition 3.9 and Corollary 3.10, every unitary dual functor $\vee_{\pi} : \mathcal{C} \rightarrow \mathcal{C}^{\text{mop}}$ on a unitary multitensor category \mathcal{C} gives a bi-involutive structure $(\bar{\cdot}^{\pi}, \nu^{\pi}, \varphi^{\pi})$ as follows. On objects, we define $\bar{c}^{\pi} := c^{\vee_{\pi}}$, and on morphisms $f \in \mathcal{C}(a \rightarrow b)$, we define

$$\bar{f}^{\pi} := \begin{array}{c} \bar{a}^{\pi} \\ \downarrow \\ \boxed{(\text{coev}_a^{\pi})^{\dagger}} \\ \downarrow \\ a \\ \downarrow \\ \boxed{f^{\dagger}} \\ \downarrow \\ b \\ \downarrow \\ \boxed{(\text{ev}_b^{\pi})^{\dagger}} \\ \downarrow \\ \bar{b}^{\pi} \end{array} = f^{\vee_{\pi} \dagger} = f^{\dagger \vee_{\pi}}.$$

We take the canonical unitary monoidal natural isomorphisms $\nu^{\pi} = \{\nu_{a,b}^{\pi} : \bar{a}^{\pi} \otimes \bar{b}^{\pi} \rightarrow \overline{b \otimes a}^{\pi}\}$ and $\varphi^{\pi} : \text{id} \Rightarrow \bar{\bar{\cdot}}^{\pi}$ induced by \vee_{π} .

Remark 3.39. Notice that the data of a bi-involutive structure $(\bar{\cdot}^{\pi}, \nu^{\pi}, \varphi^{\pi})$ is weaker than that of the dual functor \vee_{π} as we have *forgotten* the evaluation and coevaluation morphisms $\text{ev}_c^{\pi}, \text{coev}_c^{\pi}$ for $c \in \mathcal{C}$. Thus we *cannot* recover \vee_{π} from $(\bar{\cdot}^{\pi}, \nu^{\pi}, \varphi^{\pi})$.

In order to discuss unitary equivalence of bi-involutive structures, we first define the notion of a bi-involutive tensor functor from [HP17].

Definition 3.40. A *bi-involutive tensor functor* between bi-involutive tensor \mathbf{C}^* categories

$$(F, \mu, \chi) : (\mathcal{C}, \bar{\cdot}^{\mathcal{C}}, \nu^{\mathcal{C}}, \varphi^{\mathcal{C}}) \rightarrow (\mathcal{D}, \bar{\cdot}^{\mathcal{D}}, \nu^{\mathcal{D}}, \varphi^{\mathcal{D}})$$

is a dagger tensor functor (F, μ) (our convention is $\mu_{a,b} : F(a) \otimes F(b) \rightarrow F(a \otimes b)$) equipped with a unitary natural isomorphism $\chi_c : F(\bar{c}) \rightarrow \overline{F(c)}$ which is monoidal with respect to $\mu, \nu^{\mathcal{C}}, \nu^{\mathcal{D}}$ and involutive with respect to $\varphi^{\mathcal{C}}, \varphi^{\mathcal{D}}$. That is, the following diagrams commute:

$$\begin{array}{ccc} F(\bar{a}) \otimes F(\bar{b}) & \xrightarrow{\mu_{\bar{a}, \bar{b}}} & F(\bar{a} \otimes \bar{b}) \xrightarrow{F(\nu_{a,b}^{\mathcal{C}})} & F(\overline{b \otimes a}) & & F(a) \xrightarrow{F(\varphi_a^{\mathcal{C}})} & F(\bar{a}) \\ \downarrow \chi_a \otimes \chi_b & & & \downarrow \chi_{b \otimes a} & & \downarrow \varphi_{F(a)}^{\mathcal{D}} & \downarrow \chi_{\bar{a}} \\ \overline{F(a)} \otimes \overline{F(b)} & \xrightarrow{\nu_{F(a), F(b)}^{\mathcal{D}}} & \overline{F(b) \otimes F(a)} \xrightarrow{\mu_{b,a}} & \overline{F(b \otimes a)} & & \overline{F(a)} \xleftarrow{\chi_a} & \overline{F(\bar{a})} \end{array}$$

¹²Using the notation of [DSPS13], \mathcal{C}^{mp} denotes the tensor category obtained from \mathcal{C} by reversing the order of tensor product. In other words, $(\bar{\cdot}, \nu) : \mathcal{C} \rightarrow \mathcal{C}$ is conjugate-linear and anti-monoidal.

¹³Monoidality is similar to (4).

Definition 3.41. Two bi-involutive structures $(\bar{\cdot}^i, \nu^i, \varphi^i)$ on \mathcal{C} for $i = 1, 2$ are *unitarily equivalent* if there is an anti-monoidal involutive unitary natural isomorphism $\chi : \bar{\cdot}^1 \Rightarrow \bar{\cdot}^2$. This means that χ satisfies the commutative diagrams in Definition 3.40 substituting \mathcal{D} with \mathcal{C} and $F : \mathcal{C} \rightarrow \mathcal{D}$ with $\text{id}_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}$.

Remark 3.42. Given a bi-involutive structure $(\bar{\cdot}, \nu, \varphi)$ on a unitary multitensor category \mathcal{C} , an autoequivalence $\chi \in \text{Aut}((\bar{\cdot}, \nu, \varphi))$ consists of a unitary $\chi_c \in \mathcal{C}(\bar{c} \rightarrow \bar{c})$ for all $c \in \mathcal{C}$ such that for all $a, b \in \mathcal{C}$, $\chi_{b \otimes a} \circ \nu_{a,b} = \nu_{a,b} \circ (\chi_a \otimes \chi_b)$ and $\overline{\chi_a} = \chi_{\bar{a}}^{-1}$. Similar to Remark 2.5 and Lemma 3.25, looking at simple objects, we get a canonical isomorphism between $\text{Aut}((\bar{\cdot}, \nu, \varphi))$ and $\text{Hom}(\mathcal{U}^{\text{op}} \rightarrow U(1))$, but notice $g \mapsto g^{-1}$ gives an isomorphism $\mathcal{U} \cong \mathcal{U}^{\text{op}}$. This means for any two bi-involutive structures $(\bar{\cdot}^i, \nu^i, \varphi^i)$ on \mathcal{C} for $i = 1, 2$, $\text{Hom}((\bar{\cdot}^1, \nu^1, \varphi^1) \rightarrow (\bar{\cdot}^2, \nu^2, \varphi^2))$ is either empty or a torsor for $\text{Hom}(\mathcal{U}^{\text{op}} \rightarrow U(1))$. Hence there is not a *unique* unitary equivalence between two unitarily equivalent bi-involutive structures. However, we will see in the proof of Corollary B below that given two unitary dual functors, there is a *canonical* unitary equivalence between their induced bi-involutive structures.

Example 3.43. When (N, tr) is a II_1 factor with its canonical trace, there are two distinguished unitary dual functors that are often used in applications. One is the balanced dual functor giving the canonical spherical structure corresponding to the trivial homomorphism $\pi = 1$. The other is obtained from the grading on $\text{Bim}_{\text{bf}}(N)$ given by taking the ratio of the left to right von Neumann dimension. When $N = R$, the hyperfinite II_1 factor, this grading is faithful, since the fundamental group of R is $\mathbb{R}_{>0}$ [MvN43]. Taking the group homomorphism $\text{id} : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ as in Remark 3.33 gives the *tracial* unitary dual functor. Calculating the universal grading group of $\text{Bim}_{\text{bf}}(R)$ remains an important open question. Interestingly, both the spherical and tracial unitary dual functors induce unitarily equivalent bi-involutive structures as was noted in [JP19, Rem. 2.14].

Motivated by Example 3.43, we now prove the following.

Corollary (Corollary B). *Any two bi-involutive structures on a unitary multitensor category induced by unitary dual functors are canonically unitarily equivalent.*

Proof. Suppose \vee_1 and \vee_2 are two unitary dual functors on \mathcal{C} , and let π_1 and π_2 be the corresponding homomorphisms in $\text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$. While the unique monoidal natural isomorphism $\zeta : \vee_2 \Rightarrow \vee_1$ from (3) is *not* unitary, it can be *rescaled* as in Remark 3.35 to obtain a canonical unitary equivalence between the bi-involutive structures induced by \vee_1 and \vee_2 . Indeed, we define for a simple $c \in \mathcal{C}$ with $\text{gr}(c) = g$,

$$\chi_c := \left(\frac{\pi_2(g)}{\pi_1(g)} \right)^{1/4} c^{\vee_2} \text{ (cup) } c^{\vee_1} = \left(\frac{\pi_2(g)}{\pi_1(g)} \right)^{1/4} \zeta_c.$$

The above formula is derived as follows. First, we rescale \vee_1 and \vee_2 to get balanced dual functors \vee_1^b and \vee_2^b as in (24). By Corollary 3.32, the unique monoidal natural isomorphism from (3) $\zeta^b : \vee_2^b \Rightarrow \vee_1^b$ is necessarily unitary. Notice now that $\chi_c = \zeta_c^b$ as morphisms from $c^{\vee_2} = c^{\vee_2^b}$ to $c^{\vee_1} = c^{\vee_1^b}$, as the rescaling procedure fixes the dual objects. Hence χ is unitary. It is now straightforward to verify that for all $a, b \in \mathcal{C}$, $\chi_{b \otimes a} \circ \nu_{a,b} = \nu_{a,b} \circ (\chi_a \otimes \chi_b)$ and $\overline{\chi_a} = \chi_{\bar{a}}^{-1}$. \square

Remark 3.44. While the canonical unitary equivalence χ from the proof of Corollary B is not unique by Remark 3.42, it is the unique unitary natural isomorphism which can be obtained

from the unique monoidal natural isomorphism $\zeta : \vee_1 \Rightarrow \vee_2$ by scaling the ζ_c for simple $c \in \mathcal{C}$ by strictly positive real numbers. Hence if we have three unitary dual functors \vee_i for $i = 1, 2, 3$ which induce bi-involutive structures $(\bar{\cdot}^i, \nu^i, \varphi^i)$ for $i = 1, 2, 3$, then the canonical unitary equivalence χ^{12} composed with the canonical unitary equivalence χ^{23} is equal to the canonical unitary equivalence χ^{13} .

We finish this section by providing some important results on bi-involutive tensor functors.

Proposition 3.45. *Suppose $(F, \mu) : (\mathcal{C}, \vee) \rightarrow (\mathcal{D}, \vee)$ is a dagger tensor functor between unitary multitensor categories equipped with unitary dual functors, and let $\varphi^{\mathcal{C}}$ and $\varphi^{\mathcal{D}}$ be the induced unitary pivotal structures. The following are equivalent.*

- (1) (F, μ) is pivotal with respect $\varphi^{\mathcal{C}}$ and $\varphi^{\mathcal{D}}$.
- (2) The canonical isomorphism δ_c from (10) is unitary for all $c \in \mathcal{C}$.

Proof. For notational simplicity, we simply denote evaluations and coevaluations in this proof by ev and coev . Recall from (11) that (F, μ) is pivotal if and only if for all $c \in \mathcal{C}$, $\delta_c^\vee \circ \varphi_{F(c)}^{\mathcal{D}} = \delta_{c^\vee} \circ F(\varphi_c^{\mathcal{C}})$. This equality holds if and only if

$$F(\text{coev}_c^\dagger) \circ \mu_{c, c^\vee} = \text{coev}_{F(c)}^\dagger \circ (\text{id}_{F(c)} \otimes \delta_c)$$

if and only if

$$\delta_c = (\text{id}_{F(c)^\vee} \otimes F(\text{coev}_c^\dagger)) \circ (\text{id}_{F(c)^\vee} \otimes \mu_{c, c^\vee}) \circ (\text{ev}_{F(c)}^\dagger \otimes \text{id}_{F(c^\vee)})$$

if and only if $\delta_c = (\delta_c^{-1})^\dagger$ is unitary. \square

Corollary 3.46. *If either of the equivalent conditions in Proposition 3.45 hold, then (F, μ, δ) is bi-involutive.*

Proof. We must verify the diagrams in Definition 3.40 commute for (F, μ, δ) . For the first diagram, one shows both composites from $F(a^\vee) \otimes F(b^\vee) \rightarrow F(b \otimes a)^\vee$ are equal to

$$\begin{aligned} & ([F(\text{ev}_a) \circ \mu_{a^\vee, a}] \otimes \text{id}_{F(b \otimes a)^\vee}) \circ (\text{id}_{F(a^\vee)} \otimes [F(\text{ev}_b) \circ \mu_{b, b^\vee}] \otimes \text{id}_{F(a)} \otimes \text{id}_{F(b \otimes a)^\vee}) \\ & \circ (\text{id}_{F(a^\vee) \otimes F(b^\vee)} \otimes \mu_{b, a}^{-1} \otimes \text{id}_{F(b \otimes a)^\vee}) \circ (\text{id}_{F(a^\vee) \otimes F(b^\vee)} \otimes \text{coev}_{F(b \otimes a)}). \end{aligned}$$

We leave the details to the reader. For the second diagram, just notice that this is exactly the pivotality condition (11) when δ is unitary, as $\bar{\delta}_c = (\delta_c^\vee)^{-1}$. \square

The following remark is based on a suggestion of Marcel Bischoff.

Remark 3.47. Suppose that \mathcal{C}, \mathcal{D} are unitary multitensor categories which are both faithfully graded by the groupoid \mathcal{G} . Suppose that we have unitary dual functors on \mathcal{C} and \mathcal{D} , and let $(\bar{\cdot}^{\mathcal{C}}, \nu^{\mathcal{C}}, \varphi^{\mathcal{C}})$ and $(\bar{\cdot}^{\mathcal{D}}, \nu^{\mathcal{D}}, \varphi^{\mathcal{D}})$ be the induced bi-involutive structures. Suppose that $(F, \mu) : \mathcal{C} \rightarrow \mathcal{D}$ is a dagger tensor functor such that the canonical isomorphism δ_c from (10) is unitary for all $c \in \mathcal{C}$, so that (F, μ, δ) is bi-involutive by Corollary 3.46.

Suppose now that (F, μ, δ) preserves the grading groupoid \mathcal{G} , i.e., $\text{gr}(F(c)) = \text{gr}(c)$ for all homogeneous $c \in \mathcal{C}$. Picking an arbitrary $\pi \in \text{Hom}(\mathcal{G} \rightarrow \mathbb{R}_{>0})$, we can renormalize the cups and caps by π as in (23) from Remark 3.35 to get new unitary dual functors on \mathcal{C} and \mathcal{D} respectively. Note that these new dual functors need not be π -balanced unless the unitary dual functors we started with were the canonical spherical ones. However, for lack of better notation, we will denote the new evaluations and coevaluations by ev^π and coev^π respectively.

Notice that rescaling as in (23) leaves δ *unchanged!* Indeed, denoting the new canonical monoidal natural isomorphism by δ^π (again due to lack of better notation), for a homogeneous $c \in \mathcal{C}$ with $\text{gr}(c) = \text{gr}(F(c)) = g \in \mathcal{G}$, we have

$$\begin{aligned} \delta_c^\pi &= ([F(\text{ev}_c^\pi) \circ \mu_{c^\vee, c}] \otimes \text{id}_{F(c)^\vee}) \circ (\text{id}_{F(c)} \otimes \text{coev}_{F(c)}^\pi) \\ &= \pi(g)^{1/4} \pi(g)^{-1/4} \cdot (([F(\text{ev}_c) \circ \mu_{c^\vee, c}] \otimes \text{id}_{F(c)^\vee}) \circ (\text{id}_{F(c)} \otimes \text{coev}_{F(c)})) = \delta_c. \end{aligned}$$

Since the bi-involutive structures of \mathcal{C} and \mathcal{D} did not change by Corollary B, we conclude from Proposition 3.45 that (F, μ) is pivotal with respect to the new unitary pivotal structures, as the pivotal functor condition (11) still holds.

4. Planar algebras and projection categories

Planar algebras come in many variants; among them are *unoriented* [Jon11, Def. 1.1.1] (see also [MPS10]), *oriented* [Jon11, Def. 1.1.5] *disoriented* [CMW09], and *shaded* [Jon11, Def. 1.1.4] (see also [Pet10, Jon12]).

Shaded planar algebras were first defined in [Jon99] to axiomatize the standard invariant of a finite index subfactor. Since, they have been a valuable tool in the construction [BMPS12, MP15b] and classification [JMS14, AMP15] of subfactor planar algebras as they give a generators and relations approach to subfactor theory.

The following theorem is known to experts [MPS10, Gho11, Yam12, BHP12, HPT16b].

Theorem 4.1 (Folklore). *There is an equivalence of categories¹⁴*

$$\{\text{Oriented planar algebras}\} \longleftrightarrow \{\text{Pairs } (\mathcal{C}, X) \text{ with } \mathcal{C} \text{ a pivotal category and generator } X \in \mathcal{C}\}$$

Here, we call $X \in \mathcal{C}$ a generator if every object of \mathcal{C} is isomorphic to a direct summand of a direct sum of tensor powers of X and X^\vee .

Theorem 4.1 holds for many sub-classes of planar algebras and pivotal categories. We provide a helpful dictionary below:

Planar algebras	Pivotal categories with generators
unoriented	symmetrically self-dual generator
connected ($\dim(\mathcal{P}_0) = 1$)	$1_{\mathcal{C}}$ simple
2-shaded	partition $1_{\mathcal{C}} = 1_+ \oplus 1_-$ with generator $X = 1_+ \otimes X \otimes 1_-$
2-shaded connected	in addition to line above, $1_+, 1_-$ are simple
semisimple	semisimple
finite depth	finitely semisimple
spherical	spherical
\mathbb{C}^*	\mathbb{C}^* with unitary pivotal structure

For example, we get an equivalence of categories between finite depth subfactor (2-shaded connected spherical \mathbb{C}^*) planar algebras and pairs (\mathcal{C}, X) where \mathcal{C} is a finitely semisimple unitary multifusion category with its canonical spherical structure (see Corollary 3.34) such that $1_{\mathcal{C}}$ decomposes into simples as $1_{\mathcal{C}} = 1_+ \oplus 1_-$ and $X = 1_+ \otimes X \otimes 1_-$ generates \mathcal{C} .

¹⁴ Pairs (\mathcal{C}, X) form a 2-category which is equivalent to a 1-category in the following sense. Between any two 1-morphisms, there is at most one 2-morphism, which is necessarily invertible when it exists. We refer the reader to [HPT16b, Lem. 3.5] and the paragraph thereafter for more details on this subtlety.

4.1 Correspondence between planar algebras and projection categories In all of the above cases, one can recover the original pivotal category from the planar algebra as the *idempotent category*, or in the C^* cases as the *projection category*. We only spell this out here in the case of 2-shaded planar C^* algebras.

Definition 4.2. The projection category of a shaded C^* planar algebra \mathcal{P}_\bullet is the unitary multitensor category with unitary pivotal structure defined as follows:

- The objects are the projections $p \in \mathcal{P}_{n,\pm}$ ($p = p^\dagger = p^2$) and the tensor product is horizontal juxtaposition (which is zero if the two shadings do not agree).
- The morphisms spaces are given for $p \in \mathcal{P}_{n,\pm}$ and $q \in \mathcal{P}_{n',\pm'}$ by

$$\mathrm{Hom}(p \rightarrow q) = \delta_{\pm'=\pm} \delta_{n' \equiv n \pmod{2}} \left\{ x \in \mathcal{P}_{(n+n')/2,\pm} \mid x = \begin{array}{c} \begin{array}{|c} \hline n' \\ \hline \end{array} \\ \star \textcircled{q} \\ \begin{array}{|c} \hline n' \\ \hline \end{array} \\ \star \textcircled{x} \\ \begin{array}{|c} \hline n \\ \hline \end{array} \\ \star \textcircled{p} \\ \begin{array}{|c} \hline n \\ \hline \end{array} \end{array} \right\}.$$

- The adjoint is the dagger structure of the planar algebra; notice that if $x \in \mathrm{Hom}(p \rightarrow q)$, then $x^\dagger \in \mathrm{Hom}(q \rightarrow p)$.
- We get a homomorphism $\pi : \mathcal{U} \rightarrow \mathbb{R}_{>0}$ by taking the ratio of left to right traces as in (15) from Lemma 3.26. The π -balanced dual of $p \in \mathcal{P}_{n,\pm}$ is given by the the 180° -rotation of p in $\mathcal{P}_{n,\mp}$, with evaluation and co-evaluation given by using cups and caps:

$$\mathrm{coev}_p := \begin{array}{c} \begin{array}{|c} \hline n \\ \hline \end{array} \\ \star \textcircled{p} \\ \begin{array}{|c} \hline n \\ \hline \end{array} \end{array} \begin{array}{c} \begin{array}{|c} \hline n \\ \hline \end{array} \\ \textcircled{d} \\ \begin{array}{|c} \hline n \\ \hline \end{array} \end{array} \star$$

$$\mathrm{ev}_p := \begin{array}{c} \begin{array}{|c} \hline n \\ \hline \end{array} \\ \textcircled{d} \\ \begin{array}{|c} \hline n \\ \hline \end{array} \end{array} \star \star \begin{array}{c} \begin{array}{|c} \hline n \\ \hline \end{array} \\ \textcircled{p} \\ \begin{array}{|c} \hline n \\ \hline \end{array} \end{array}.$$

It is straightforward to calculate from the formula for the unitary pivotal structure in (2) of Proposition 3.9 that $\varphi_p^\pi = \mathrm{id}_p$.

The generator corresponds to the unshaded-shaded strand $\blacksquare \in \mathcal{P}_{1,+}$.

Definition 4.3. Conversely, given a unitary multitensor category \mathcal{C} where $1_{\mathcal{C}} = 1_+ \oplus 1_-$ is an orthogonal decomposition (with 1_{\pm} not necessarily simple) together with a fixed $\pi \in \mathrm{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$ and generator $X = 1_+ \otimes X \otimes 1_-$, we obtain a shaded C^* planar algebra \mathcal{P}_\bullet by a unitary version of [Gho11, §4]. Let $(X^{\vee\pi}, \mathrm{ev}_X^\pi, \mathrm{coev}_X^\pi)$ be the π -balanced dual of X as in Proposition 3.30. We use the notation

$$X^{\mathrm{alt}\otimes n} := \underbrace{X \otimes X^{\vee\pi} \otimes \dots \otimes X^?}_{n \text{ tensorands}} \quad (\overline{X}^\pi)^{\mathrm{alt}\otimes n} := \underbrace{X^{\vee\pi} \otimes X \otimes \dots \otimes (X^{\vee\pi})^?}_{n \text{ tensorands}}$$

where $X^? = X$ if n is odd and $X^{\vee\pi}$ if X is even, and $(X^{\vee\pi})^? = X$ if n is even and $X^{\vee\pi}$ if n is odd. The box spaces are defined for $n \geq 0$ by

$$\mathcal{P}_{n,+} := \mathcal{C}(X^{\mathrm{alt}\otimes n} \rightarrow X^{\mathrm{alt}\otimes n}) \quad \mathcal{P}_{n,-} := \mathcal{C}((X^{\vee\pi})^{\mathrm{alt}\otimes n} \rightarrow (X^{\vee\pi})^{\mathrm{alt}\otimes n}). \quad (25)$$

The action of tangles is via the graphical calculus for pivotal categories; details appear in [Gho11, §4]. For our purposes, we specify the actions of the following shaded planar tangles, which determines the action of every shaded planar tangle.

- The strand is the identity morphism $\blacksquare := \mathrm{id}_X$ and $\blacksquare := \mathrm{id}_{X^{\vee\pi}}$
- Caps and cups which are shaded above are given by $\frown := \mathrm{ev}_X^\pi$ and $\smile := \mathrm{coev}_X^\pi$

- Caps and cups which are shaded below are given by $\frown := (\text{coev}_X^\pi)^\dagger$ and $\smile := (\text{ev}_X^\pi)^\dagger$
- Vertical join is composition in \mathcal{C} $\begin{array}{c} \boxed{g} \\ \boxed{f} \end{array} := g \circ f$
- Horizontal join is tensor product in \mathcal{C} $\boxed{f} \boxed{g} := f \otimes g$.

The \dagger -structure on \mathcal{P}_\bullet is just \dagger on morphisms in \mathcal{C} .

4.2 Making closed loops scalar valued For this section, we assume \mathcal{C} is a unitary multi-tensor category such that $\dim(\mathcal{C}(1_{\mathcal{C}} \rightarrow 1_{\mathcal{C}})) = r$ and that \mathcal{C} is faithfully graded by the groupoid \mathcal{M}_r from Example 3.19.

While \mathcal{C} has a canonical spherical structure from Corollary 3.34, it is not always the most relevant one for planar algebraic purposes, including graph planar algebra embedding. Example 4.7 below discusses the graph planar algebra in detail. Given an object $X \in \mathcal{C}$ which partitions $1_{\mathcal{C}}$ into a source and target summand, one may desire that closed loops count for scalars in the planar algebra associated to (\mathcal{C}, \vee, X) for some unitary dual functor \vee . The conditions on \vee are exactly provided by the recent article [GL19], which introduced the notion of standard solutions for the conjugate equations for a unitary multitensor category *with respect to an object* $X \in \mathcal{C}$ in the more general context of C^* 2-categories. We now rephrase their definition in our setup.

Definition 4.4 ([GL19, Def. 7.25 and 7.29]). Suppose $X \in \mathcal{C}$ such that there is an orthogonal decomposition $1_{\mathcal{C}} = 1_+ \oplus 1_-$, which are not necessarily simple, such that $X = 1_+ \otimes X \otimes 1_-$. Let V_{\pm} be a set of representatives of the simple summands of 1_{\pm} .

Let $D_X \in M_{V_+ \times V_-}(\mathbb{C})$ be the matrix whose uv -th entry is equal to the positive spherical dimension $\dim(u \otimes X \otimes v)$ using the canonical spherical structure from Corollary 3.34. Let $d_X > 0$ such that d_X^2 is the common Frobenius-Perron eigenvalue of $D_X D_X^T$ and $D_X^T D_X$. Let λ_+ and λ_- be the Frobenius-Perron eigenvectors for $D_X D_X^T$ and $D_X^T D_X$ which are normalized so that

$$\sum_{u \in V_+} \lambda_+(u)^2 = 1 = \sum_{v \in V_-} \lambda_-(v)^2.$$

We define

$$\lambda := \begin{pmatrix} \lambda_+ \\ \lambda_- \end{pmatrix} \in \mathbb{C}^r. \quad (26)$$

The unitary dual functor corresponding to the groupoid homomorphism $\pi_X : \mathcal{M}_r \rightarrow \mathbb{R}_{>0}$ given by

$$\pi_X(E_{uv}) := \left(\frac{\lambda(u)}{\lambda(v)} \right)^2. \quad (27)$$

is called *standard with respect to* X . The unitary dual functor corresponding to the groupoid homomorphism $\pi_X^\ell : \mathcal{M}_r \rightarrow \mathbb{R}_{>0}$ given by

$$\pi_X^\ell(E_{uv}) := \begin{cases} d_X^{-1} \left(\frac{\lambda(u)}{\lambda(v)} \right)^2 & \text{if } u \in V_+ \text{ and } v \in V_- \\ d_X \left(\frac{\lambda(u)}{\lambda(v)} \right)^2 & \text{if } u \in V_- \text{ and } v \in V_+ \\ \left(\frac{\lambda(u)}{\lambda(v)} \right)^2 & \text{else.} \end{cases} \quad (28)$$

is called *lopsided with respect to* X .

The following lemma is immediate from [GL19].

Lemma 4.5.

- (1) The 2-shaded C^* planar algebra \mathcal{P}_\bullet corresponding to the triple $(\mathcal{C}, X, \vee_{\pi_X})$ with $\text{id}_X = \blacksquare \in \mathcal{P}_{1,+}$ satisfies

$$\textcircled{\text{lightblue}} = d_X \text{id}_{1_+} \quad \textcircled{\text{lightblue}} \textcircled{\text{white}} = d_X \text{id}_{1_-}. \quad (29)$$

- (2) The 2-shaded C^* planar algebra \mathcal{P}_\bullet^ℓ corresponding to the triple $(\mathcal{C}, X, \vee_{\pi_X^\ell})$ with $\text{id}_X = \blacksquare \in \mathcal{P}_{1,+}^\ell$ satisfies

$$\textcircled{\text{lightblue}} = \text{id}_{1_+} \quad \textcircled{\text{lightblue}} \textcircled{\text{white}} = d_X^2 \text{id}_{1_-}. \quad (30)$$

Remark 4.6. The lopsided planar algebra is obtained from the standard planar algebra as in Remark 3.47 by replacing the standard ev_X and coev_X by $d_X^{1/2} \text{ev}_X$ and $d_X^{-1/2} \text{coev}_X$. Notice this lopsided unitary pivotal structure varies slightly from the *non-unitary* lopsided convention from [MP14, §1.1], which replaces ev_X and coev_X by $d_X \text{ev}_X$ and $d_X^{-1} \text{coev}_X$, but does *not* scale ev_X^\dagger nor coev_X^\dagger . Often, it is computationally simpler to calculate a graph planar algebra embedding with respect to the non-unitary lopsided pivotal structure, as the number fields are more manageable. It is still the case that non-unitary lopsided embeddings give standard unitary embeddings by [MP14].

Example 4.7. Let $\Gamma = (V_+, V_-, E)$ be a finite connected bipartite graph with even/+ vertices V_+ , odd/- vertices V_- , and edges E . We consider an edge $\varepsilon \in E$ as directed from + to - with source $s(\varepsilon) \in V_+$ and target $t(\varepsilon) \in V_-$. We write ε^* for the same edge with the opposite direction. Let λ denote any Frobenius-Perron eigenvector of the adjacency matrix of Γ .

Denote by n_\pm the number of vertices in V_\pm . Let $\mathcal{M}_\pm = \text{Hilb}^{n_\pm}$, where Hilb is the category of finite dimensional Hilbert spaces considered as a semisimple C^* category. We pick distinguished simples of \mathcal{M}_\pm which we name by the vertices in V_\pm . Define $\mathcal{M} = \mathcal{M}_+ \oplus \mathcal{M}_-$, which consists of one copy of Hilb for every vertex of Γ .

Now consider $\text{End}^\dagger(\mathcal{M})$, which we identify with the unitary multifusion category $\text{Mat}_{n \times n}^\dagger$. The simple objects are the $E_{u,v}$ for $u, v \in V_+ \amalg V_-$ with fusion rule $E_{u,v} \otimes E_{w,x} = \delta_{v=w} E_{u,x}$, and $1 = \bigoplus_{v \in V_+ \amalg V_-} E_{v,v}$. It is straightforward to verify that the unique spherical structure from Corollary 3.34 is given by $\text{ev}_{E_{u,v}} : E_{v,u} \otimes E_{u,v} = E_{v,v} \hookrightarrow 1$ and $\text{coev}_{E_{u,v}} : 1 \rightarrow E_{u,u} = E_{u,v} \otimes E_{v,u}$. Notice that the canonical spherical structure φ satisfies $\varphi_{E_{u,v}} = \text{id}_{E_{u,v}}$ for all vertices u, v .

Observe the universal grading groupoid \mathcal{U} of $\text{End}^\dagger(\mathcal{M})$ is $\mathcal{M}_{n_+ + n_-}$. By Corollary 3.32, we get a canonical π -balanced unitary dual functor from the homomorphism $\pi : \mathcal{M}_{n_+ + n_-} \rightarrow \mathbb{R}_{>0}$ given by (27). By direct computation as in the proof of Proposition 3.30, $\text{ev}_{E_{u,v}}^\pi$ and $\text{coev}_{E_{u,v}}^\pi$ are given by renormalizing the canonical 1-balanced evaluation and coevaluation maps:

$$\text{ev}_{E_{u,v}}^\pi := \left(\frac{\lambda(u)}{\lambda(v)} \right)^{1/2} \text{ev}_{E_{u,v}} \quad \text{coev}_{E_{u,v}}^\pi := \left(\frac{\lambda(v)}{\lambda(u)} \right)^{1/2} \text{coev}_{E_{u,v}}.$$

Thus $\dim_L^\pi(E_{u,v}) = \frac{\lambda(u)}{\lambda(v)}$ and $\dim_R^\pi(E_{u,v}) = \frac{\lambda(v)}{\lambda(u)}$.

We now pick a distinguished dagger functor $F \in \text{End}^\dagger(\mathcal{M})$ together with its π -balanced dual

$$F = \bigoplus_{\varepsilon \in E} E_{s(\varepsilon), t(\varepsilon)} \quad F^{\vee\pi} = \bigoplus_{\varepsilon \in E} E_{t(\varepsilon), s(\varepsilon)}. \quad (31)$$

Since Γ is connected, we see that F generates $\text{End}^\dagger(\mathcal{M})$. Define $1_+ := \bigoplus_{u \in V_+} E_{u,u}$ and $1_- := \bigoplus_{v \in V_-} E_{v,v}$, and note that $1 = 1_+ \oplus 1_-$ is an orthogonal decomposition of the unit object such that $F = 1_+ \otimes F \otimes 1_-$.

Let \mathcal{G}_\bullet be the corresponding shaded planar C^* algebra corresponding to $(\text{End}^\dagger(\mathcal{M}), F, \vee_\pi)$ under Theorem 4.1, which was described in Definition 4.3. We may identify $\mathcal{G}_{n,\pm}$ defined as in (25) as the complex vector space whose basis consists of the loops of length $2n$ on Γ starting at a \pm vertex in V_\pm . For example,

$$\begin{aligned} \mathcal{G}_{n,+} &:= \text{Hom}_{\text{End}^\dagger(\mathcal{M})}(F^{\text{alt}\otimes n} \rightarrow F^{\text{alt}\otimes n}) \\ &\cong \text{Hom}_{\text{End}^\dagger(\mathcal{M})}(1 \rightarrow (F \otimes F^{\vee\pi})^{\otimes n}) \\ &\cong \bigoplus_{v \in V_+} \bigoplus_{\varepsilon_1, \dots, \varepsilon_{2n} \in E} \text{Hom}_{\text{End}^\dagger(\mathcal{M})}(E_{v,v} \rightarrow \\ &\qquad\qquad\qquad E_{s(\varepsilon_1),t(\varepsilon_1)} \otimes E_{t(\varepsilon_2),s(\varepsilon_2)} \otimes \cdots \otimes E_{s(\varepsilon_{2n-1}),t(\varepsilon_{2n-1})} \otimes E_{t(\varepsilon_{2n}),s(\varepsilon_{2n})}) \\ &\cong \bigoplus_{v \in V_+} \text{span}_{\mathbb{C}}\{\text{loops of length } 2n \text{ based at } v\}. \\ &\cong \text{span}_{\mathbb{C}}\{\text{loops of length } 2n \text{ based at an even/+ vertex}\}. \end{aligned}$$

Under this identification, it is straightforward to verify that the actions of the shaded planar tangles described in Definition 4.3 exactly correspond to the actions of the shaded planar tangles for the planar algebra of the bipartite graph Γ from [Jon00].

From Theorem 4.1 and the discussion in Example 4.7, we get the following.

Proposition 4.8. *Under the equivalence of categories in Theorem 4.1 for shaded C^* planar algebras, the bipartite graph planar algebra \mathcal{G}_\bullet corresponds to the unitary multifusion category $\text{End}^\dagger(\mathcal{M})$ with (non-spherical!) unitary pivotal structure obtained from the standard groupoid homomorphism (27) with respect to F_Γ as defined in (31).*

5. Spherical states on planar algebras and multitensor categories

Motivated by the example of the graph planar algebra, we now define the notion of a spherical state with respect to a partition on a unitary multitensor category. Such a spherical state can be chosen to ‘correct’ for a non-spherical unitary pivotal structure on a unitary multitensor category if the corresponding $\pi \in \text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0})$ actually comes from a faithful grading by the groupoid \mathcal{M}_r from Example 3.19.

5.1 Evaluable planar algebras and spherical states Below, we discuss sphericity and evaluability for *semisimple* shaded planar algebras, i.e., each $\mathcal{P}_{n,\pm}$ is a finite dimensional complex semisimple algebra under the usual multiplication in \mathcal{P}_\bullet .

Definition 5.1. A shaded planar algebra is called *evaluative* if $\dim(\mathcal{P}_{0,\pm}) = 1$. An evaluative shaded planar algebra is called *spherical* if for all $x \in \mathcal{P}_{1,+}$ the following two scalars in $\mathcal{P}_{0,\pm} \cong \mathbb{C}$ (via mapping the empty diagrams to $1_{\mathbb{C}}$) agree:

For non-evaluative shaded planar algebras, [Jon11] defines sphericity in terms of a *measure* on \mathcal{P}_\bullet , which is a pair of linear functionals ψ_\pm on the finite dimensional abelian complex semisimple algebras $\mathcal{P}_{0,\pm}$. A measure (ψ_+, ψ_-) is called:

- a *state* if $\psi_{\pm}(p) \geq 0$ for every projection $p \in \mathcal{P}_{0,\pm}$,
- a *faithful state* if $\psi_{\pm}(p) > 0$ for every non-zero projection $p \in \mathcal{P}_{0,\pm}$, and
- *spherical* if for all $x \in \mathcal{P}_{1,+}$,

$$\psi_- \left(\left(\begin{array}{c} \text{---} \\ \text{---} \end{array} \right) \begin{array}{c} \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} \right) = \psi_+ \left(\left(\begin{array}{c} \text{---} \\ \text{---} \end{array} \right) \begin{array}{c} \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} \right).$$

Example 5.2. The graph planar algebra is in general not spherical. For example, taking any edge ε which connects two vertices of distinct weights, the projection $[\varepsilon\varepsilon^*] \in \mathcal{G}_{1,+}$ has distinct left and right traces. However, if we normalize the Frobenius-Perron eigenvector λ so that $\sum_{u \in V_+} \lambda(u)^2 = 1 = \sum_{v \in V_-} \lambda(v)^2$, then $\psi(p_v) := \lambda(v)^2$ defines a spherical faithful state on \mathcal{G}_{\bullet} [Jon00, Prop. 3.4].

Remark 5.3. An evaluable shaded planar algebra is spherical if and only if its pivotal projection multitensor category is spherical. We will define the concepts of measure and (spherical faithful) state for a multitensor category in Section 5.2 below.

Remark 5.4 ([Jon01, §8]). If \mathcal{P}_{\bullet} is a shaded planar (\dagger -)algebra with a spherical faithful state, then any evaluable planar (\dagger -)subalgebra $\mathcal{Q}_{\bullet} \subset \mathcal{P}_{\bullet}$ is spherical. If in addition \mathcal{P}_{\bullet} is \mathbb{C}^* with finite dimensional box spaces, then any evaluable $\mathcal{Q}_{\bullet} \subset \mathcal{P}_{\bullet}$ is a subfactor planar algebra.

5.2 Evaluable multitensor subcategories and spherical states With the graph planar algebra in mind, we now define the notions of measure and state *with respect to a partition* Π on a pivotal multitensor category which generalizes the similar notion for shaded planar algebras from §5.1.

Definition 5.5. Given a multitensor category \mathcal{C} , a *partition* Π of $1_{\mathcal{C}}$ consists of a family of mutually orthogonal projections $\{p\} \subset \text{End}_{\mathcal{C}}(1_{\mathcal{C}})$ such that $\sum_{p \in \Pi} p = 1$.

Given a multitensor category \mathcal{C} with a partition Π of $1_{\mathcal{C}}$, a *measure with respect to* Π is a linear functional ψ on the finite dimensional abelian semisimple \mathbb{C} -algebra $\text{End}_{\mathcal{C}}(1_{\mathcal{C}})$. We call a measure ψ with respect to Π :

- a *state* if $\psi(p) \geq 0$ for every idempotent $p \in \text{End}_{\mathcal{C}}(1_{\mathcal{C}})$ and $\psi(p) = 1$ for all $p \in \Pi$.
- a *faithful state* if ψ is a state and $\psi(p) > 0$ for every non-zero idempotent $p \in \text{End}_{\mathcal{C}}(1_{\mathcal{C}})$. (Note that if \mathcal{C} has simple tensor unit, there is a unique state, which is automatically faithful.)
- *spherical* if (\mathcal{C}, φ) is pivotal and $\psi \circ \text{tr}_L = \psi \circ \text{tr}_R$ for all $c \in \mathcal{C}$.

When $\Pi = \{\text{id}_{1_{\mathcal{C}}}\}$ is the trivial partition of $1_{\mathcal{C}}$, we omit Π from the notation and simply refer to measures and (spherical faithful) states.

Remark 5.6. Note that in the case where (\mathcal{C}, φ) has non-simple tensor unit, having a faithful spherical state is additional structure over being pivotal. However, if (\mathcal{C}, φ) is spherical, then setting Π to be the set of all minimal projections in $\text{End}_{\mathcal{C}}(1_{\mathcal{C}})$, there is a canonical faithful spherical state with respect to Π given by $\psi(p) = 1$ for every $p \in \Pi$.

Example 5.7. By Example 5.2, the unitary multifusion category of projections of the graph planar algebra has a spherical faithful state with respect to the induced unitary dual functor and the partition $\Pi := \{\sum_{u \in V_+} p_u, \sum_{v \in V_-} p_v\}$.

Definition 5.8. Suppose \mathcal{C} is a multitensor category with partition Π of $1_{\mathcal{C}}$. For $p \in \Pi$, denote by 1_p the summand of $1_{\mathcal{C}}$ corresponding to $p \in \text{End}_{\mathcal{C}}(1_{\mathcal{C}})$. We call a unital multitensor subcategory $\mathcal{D} \subset \mathcal{C}$ *evaluable with respect to Π* if for every $p \in \Pi$, $1_p \in \mathcal{D}$ and $\text{End}_{\mathcal{D}}(1_p) = \mathbb{C}p$, i.e., $1_{\mathcal{D}} = \bigoplus_{p \in \Pi} 1_p$ exhibits the decomposition of $1_{\mathcal{D}}$ into simple objects of \mathcal{D} .

Proposition 5.9. *Let (\mathcal{C}, φ) be a pivotal multitensor category with Π a partition of $1_{\mathcal{C}}$. Suppose ψ is a spherical state on $\text{End}_{\mathcal{C}}(1_{\mathcal{C}})$ with respect to Π . Then if \mathcal{D} is a unital multitensor subcategory of \mathcal{C} such that $\varphi_d \in \mathcal{D}(d \rightarrow d)$ for all $d \in \mathcal{D}$ and which is evaluable with respect to Π , then $(\mathcal{D}, \varphi|_{\mathcal{D}})$ is spherical.*

Proof. Since $\varphi_d \in \mathcal{D}(d \rightarrow d)$ for all $d \in \mathcal{D}$, we have $\text{tr}_L^{\mathcal{C}} = \text{tr}_L^{\mathcal{D}}$ and $\text{tr}_R^{\mathcal{C}} = \text{tr}_R^{\mathcal{D}}$ for all $d \in \mathcal{D}$. Suppose $f \in \mathcal{D}(c \rightarrow d)$. Since \mathcal{D} is evaluable with respect to Π , $\Pi \subset \text{End}_{\mathcal{D}}(1_{\mathcal{D}})$ is the set of minimal projections (which may not be minimal in $\text{End}_{\mathcal{C}}(1_{\mathcal{C}})$). Then for any $p, q \in \Pi$,

$$\text{tr}_L(p \otimes f \otimes q) = \psi(\text{tr}_L(p \otimes f \otimes q))p \quad \text{tr}_R(p \otimes f \otimes q) = \psi(\text{tr}_R(p \otimes f \otimes q))q$$

since f is a morphism in \mathcal{D} , and \mathcal{D} is evaluable with respect to Π . Now since ψ is a spherical state on $\text{End}_{\mathcal{C}}(1_{\mathcal{C}})$, we have $\psi(\text{tr}_L(p \otimes f \otimes q)) = \psi(\text{tr}_R(p \otimes f \otimes q))$, and thus \mathcal{D} is spherical. \square

Example 5.10. As we saw in §4.2, one important way that partitions Π of $1_{\mathcal{C}}$ arise naturally is from picking a distinguished object X in a unitary multitensor category whose source and range summands of $1_{\mathcal{C}}$ are orthogonal. That is $1_{\mathcal{C}} = 1_+ \oplus 1_-$ with 1_{\pm} not necessarily simple such that $X = 1_+ \otimes X \otimes 1_-$. We then set $\Pi = \{p_+, p_-\}$ where $p_{\pm} \in \mathcal{C}(1_{\mathcal{C}} \rightarrow 1_{\mathcal{C}})$ is the orthogonal projection corresponding to the summand 1_{\pm} .

Example 5.11. Building on Definition 4.4 and Examples 5.2 and 5.10, if $X = 1_+ \otimes X \otimes 1_-$ generates \mathcal{C} such that $1_{\mathcal{C}} = 1_+ \oplus 1_-$ and \mathcal{C} is faithfully graded by \mathcal{M}_r , then defining $\psi(v) := \lambda(v)^2$ for all simple summands $v \subset 1_{\mathcal{C}}$ where λ is defined as in (26) gives a spherical faithful state on $(\mathcal{C}, \vee_{\text{standard}})$, where \vee_{standard} is the standard unitary dual functor on \mathcal{C} with respect to X from (27). One calculates that for all summands $u \subset 1_-$ and $v \subset 1_+$,

$$\psi(\text{tr}_L^{\pi}(\text{id}_{u \otimes X \otimes v})) = \lambda(u)\lambda(v)(D_X)_{u,v} = \psi(\text{tr}_R^{\pi}(\text{id}_{u \otimes X \otimes v})).$$

As this equation is identical to [GL19, (7.9)], we have that their canonical left/right states of X on $\mathcal{C}(X \rightarrow X)$ are given by $\omega_{\ell} = \psi \circ \text{tr}_L^{\pi}$ and $\omega_r = \psi \circ \text{tr}_R^{\pi}$.

5.3 The spherical state correction for non-balanced unitary dual functors For this section, \mathcal{C} is a unitary multitensor category which is faithfully graded by the groupoid \mathcal{M}_r consisting of the groupoid with r objects and exactly one isomorphism between any two objects, which we can identify with the standard system of matrix units $\{E_{ij}\}$ for $M_r(\mathbb{C})$. For notational simplicity, we write \mathcal{C}_{ij} for $\mathcal{C}_{E_{ij}}$. Moreover, we assume $1_{\mathcal{C}} = \bigoplus_{i=1}^r 1_i$ is an orthogonal decomposition into simples, and we let $p_i \in \mathcal{C}(1_{\mathcal{C}} \rightarrow 1_{\mathcal{C}})$ be the minimal projection corresponding to the summand 1_i . We assume \mathcal{C} has the trivial partition $\Pi = \{\text{id}_{1_{\mathcal{C}}}\}$.

We now show that one can ‘correct’ for a unitary dual functor on \mathcal{C} which is not balanced, but comes from a $\pi \in \text{Hom}(\mathcal{M}_r \rightarrow \mathbb{R}_{>0})$. This can be viewed as a generalization of [Jon00, Thm. 3.1] for the bipartite graph planar algebra (see also Example 5.2) and [GL19, Thm. 7.39].

Example 5.12. Suppose \mathcal{C} is a (unitary) $r \times r$ multifusion category, and let $\pi : \mathcal{U} \rightarrow \mathbb{C}^{\times}$ be a homomorphism. For every subgroup $\mathcal{H} \subseteq \mathcal{U}$ corresponding to the automorphisms of a

single object, we must have $\pi(\mathcal{H}) \subset U(1)$. When \mathcal{C} is unitary and $\pi : \mathcal{U} \rightarrow \mathbb{R}_{>0}$, we have $\pi(\mathcal{H}) \subseteq U(1) \cap \mathbb{R}_{>0} = \{1\}$. Thus the canonical groupoid surjection $\mathcal{U} \twoheadrightarrow \mathcal{M}_r$ from Remark 3.17 induces an isomorphism $\text{Hom}(\mathcal{U} \rightarrow \mathbb{R}_{>0}) \cong \text{Hom}(\mathcal{M}_r \rightarrow \mathbb{R}_{>0})$.

Fact 5.13. *An element $\pi \in \text{Hom}(\mathcal{M}_r \rightarrow \mathbb{R}_{>0})$ is completely determined by its values on $E_{i+1,i}$ for $i = 1, \dots, r-1$, which can be arbitrary. Hence $\pi \mapsto (\pi(E_{i+1,i}))_{i=1}^{r-1}$ is a bijection $\text{Hom}(\mathcal{M}_r \rightarrow \mathbb{R}_{>0}) \cong \mathbb{R}_{>0}^{r-1}$.*

Lemma 5.14. *The function*

$$\psi \mapsto \left(\frac{\psi(p_{i+1})}{\psi(p_i)} \right)_{i=1}^{r-1}$$

is a bijection between faithful states ψ on \mathcal{C} with respect to Π and $\mathbb{R}_{>0}^{r-1}$.

Proof. First, consider the set \mathcal{F} of all linear functionals $\psi : \mathcal{C}(1_{\mathcal{C}} \rightarrow 1_{\mathcal{C}}) \rightarrow \mathbb{C}$ such that $\psi(p_i) \neq 0$ for all $i = 1, \dots, r$. The map $\mathcal{F} \ni \psi \mapsto \left(\frac{\psi(p_{i+1})}{\psi(p_i)} \right)_{i=1}^{r-1} \in \mathbb{R}_{>0}^{r-1}$ is clearly well defined and surjective. Moreover, we see that ψ_1 and ψ_2 map to the same element of $\mathbb{R}_{>0}^{r-1}$ if and only if they are proportional:

$$\frac{\psi_1(p_{i+1})}{\psi_1(p_i)} = \frac{\psi_2(p_{i+1})}{\psi_2(p_i)} \quad \forall i = 1, \dots, r-1 \quad \iff \quad \frac{\psi_2(p_i)}{\psi_1(p_i)} = \frac{\psi_2(p_{i+1})}{\psi_1(p_{i+1})} \quad \forall i = 1, \dots, r-1.$$

Finally, given a $\psi \in \mathcal{F}$, there is exactly one faithful state with respect to Π which is proportional to it. \square

Theorem (Theorem D). *Given a $\pi \in \text{Hom}(\mathcal{M}_r \rightarrow \mathbb{R}_{>0})$, there is a unique spherical faithful state ψ^π with respect to Π for $(\mathcal{C}, \vee_\pi, \nu^\pi, \varphi^\pi)$.*

Proof.

Step 1: Suppose $c \in \mathcal{C}_{ij}$ is simple. Then

$$\psi^\pi(\text{tr}_L^\pi(\text{id}_c)) = \psi^\pi(\text{tr}_R^\pi(\text{id}_c)) \quad \iff \quad \psi^\pi(p_j) \dim_L^\pi(c) = \psi^\pi(p_i) \dim_R^\pi(c),$$

which is equivalent to

$$\frac{\psi^\pi(p_i)}{\psi^\pi(p_j)} = \pi(E_{ij}) \stackrel{(15)}{=} \frac{\dim_L^\pi(c)}{\dim_R^\pi(c)}.$$

Notice that both π and $E_{ij} \mapsto \frac{\psi^\pi(p_i)}{\psi^\pi(p_j)}$ are groupoid homomorphisms, so the above equality holds for all simple $c \in \mathcal{C}_{ij}$ for all $i, j = 1, \dots, r$ if and only if it holds for all simple $c \in \mathcal{C}_{i+1,i}$ for all $i = 1, \dots, r-1$. This is equivalent to both homomorphisms corresponding to the same element of $\mathbb{R}_{>0}^{r-1}$ under the bijections from Fact 5.13 and Lemma 5.14 respectively. Hence there is a unique choice of ψ^π which works.

Step 2: Suppose $c \in \mathcal{C}$ is an orthogonal direct sum of n objects isomorphic to the simple object $a \in \mathcal{C}$ and $f \in \mathcal{C}(a \rightarrow a)$. Pick n isometries $v_1, \dots, v_n \in \mathcal{C}(a \rightarrow c)$ with orthogonal ranges so that $\sum_{i=1}^n v_i \circ v_i^\dagger = \text{id}_c$. Then for ψ^π defined in Step 1,

$$\begin{aligned} \psi^\pi(\text{tr}_L^\pi(f)) &= \sum_{i=1}^n \psi^\pi(\text{tr}_L^\pi(v_i \circ v_i^\dagger \circ f)) = \sum_{i=1}^n \psi^\pi(\text{tr}_L^\pi(v_i^\dagger \circ f \circ v_i)) \\ &= \sum_{i=1}^n \psi^\pi(\text{tr}_R^\pi(v_i^\dagger \circ f \circ v_i)) = \sum_{i=1}^n \psi^\pi(\text{tr}_L^\pi(v_i \circ v_i^\dagger \circ f)) = \psi^\pi(\text{tr}_R^\pi(f)). \end{aligned}$$

Step 3: Suppose $c \in \mathcal{C}$ and $f \in \mathcal{C}(c \rightarrow c)$ are arbitrary. Decompose c into an orthogonal direct sum of isotypic components and apply Step 2. \square

Acknowledgements

The author would like to thank André Henriques, Corey Jones, and Noah Snyder for many helpful conversations without which this article would not have been possible. The author would also like to thank Marcel Bischoff, Ian Charlesworth, Sam Evington, Luca Giorgetti, and Cris Negron, a.k.a. the ‘Unitary Group’ from the 2018 AMS MRC on Quantum Symmetries: Subfactors and Fusion Categories, supported by NSF DMS grant 1641020, for an extremely helpful and clarifying week of discussion on related topics. The author was supported by NSF DMS grants 1500387/1655912 and 1654159.

References

- [AH99] Marta Asaeda and Uffe Haagerup, *Exotic subfactors of finite depth with Jones indices $(5 + \sqrt{13})/2$ and $(5 + \sqrt{17})/2$* , *Comm. Math. Phys.* **202** (1999), no. 1, 1–63, [MR1686551](#), [DOI:10.1007/s002200050574](#), [arXiv:math.OA/9803044](#).
- [AMP15] Narjess Afzaly, Scott Morrison, and David Penneys, *The classification of subfactors with index at most $5\frac{1}{4}$* , 2015, [arXiv:1509.00038](#), to appear *Mem. Amer. Math. Soc.*
- [AP17] Claire Anantharaman and Sorin Popa, *An introduction to II_1 factors*, 2017, preprint available at <http://www.math.ucla.edu/~popa/books.html>.
- [BDH14] Arthur Bartels, Christopher L. Douglas, and André Henriques, *Dualizability and index of subfactors*, *Quantum Topol.* **5** (2014), no. 3, 289–345, [MR3342166](#) [DOI:10.4171/QT/53](#) [arXiv:1110.5671](#).
- [BHP12] Arnaud Brothier, Michael Hartglass, and David Penneys, *Rigid C^* -tensor categories of bimodules over interpolated free group factors*, *J. Math. Phys.* **53** (2012), no. 12, 123525, 43, [MR3405915](#) [DOI:10.1063/1.4769178](#) [arXiv:1208.5505](#).
- [Bis97] Dietmar Bisch, *Bimodules, higher relative commutants and the fusion algebra associated to a subfactor*, *Operator algebras and their applications (Waterloo, ON, 1994/1995)*, 13–63, *Fields Inst. Commun.*, 13, Amer. Math. Soc., Providence, RI, 1997, [MR1424954](#), (preview at [google books](#)).
- [BKLR15] Marcel Bischoff, Yasuyuki Kawahigashi, Roberto Longo, and Karl-Henning Rehren, *Tensor categories and endomorphisms of von Neumann algebras—with applications to quantum field theory*, *SpringerBriefs in Mathematical Physics*, vol. 3, Springer, Cham, 2015, [MR3308880](#) [DOI:10.1007/978-3-319-14301-9](#).
- [BMPS12] Stephen Bigelow, Scott Morrison, Emily Peters, and Noah Snyder, *Constructing the extended Haagerup planar algebra*, *Acta Math.* **209** (2012), no. 1, 29–82, [MR2979509](#), [arXiv:0909.4099](#), [DOI:10.1007/s11511-012-0081-7](#).
- [CHPS18] Desmond Coles, Peter Huston, David Penneys, and Srivatsa Srinivas, *The module embedding theorem via towers of algebras*, 2018, [arXiv:1810.07049](#).
- [CMW09] David Clark, Scott Morrison, and Kevin Walker, *Fixing the functoriality of Khovanov homology*, *Geom. Topol.* **13** (2009), no. 3, 1499–1582, [MR2496052](#) [DOI:10.2140/gt.2009.13.1499](#) [arXiv:math.GT/0701339](#).
- [DSPS13] Chris Douglas, Chris Schommer-Pries, and Noah Snyder, *Dualizable tensor categories*, 2013, [arXiv:1312.7188](#).
- [Egg11] J. M. Egger, *On involutive monoidal categories*, *Theory Appl. Categ.* **25** (2011), No. 14, 368–393, [MR2861112](#).
- [EGNO15] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik, *Tensor categories*, *Mathematical Surveys and Monographs*, vol. 205, American Mathematical Society, Providence, RI, 2015, [MR3242743](#) [DOI:10.1090/surv/205](#).
- [Gho11] Shamindra Kumar Ghosh, *Planar algebras: a category theoretic point of view*, *J. Algebra* **339** (2011), 27–54, [MR2811311](#), [arXiv:0810.4186](#), [DOI:10.1016/j.jalgebra.2011.04.017](#).
- [GL19] Luca Giorgetti and Roberto Longo, *Minimal index and dimension for $2-C^*$ -categories with finite-dimensional centers*, *Comm. Math. Phys.* **370** (2019), no. 2, 719–757, [MR3994584](#) [DOI:10.1007/s00220-018-3266-x](#) [arXiv:1805.09234](#).

- [GLR85] P. Ghez, R. Lima, and J. E. Roberts, *W*-categories*, Pacific J. Math. **120** (1985), no. 1, 79–109, MR808930.
- [GMP⁺18] Pinhas Grossman, Scott Morrison, David Penneys, Emily Peters, and Noah Snyder, *The Extended Haagerup fusion categories*, 2018, arXiv:1810.06076.
- [Got10] Satoshi Goto, *On Ocneanu’s theory of double triangle algebras for subfactors and classification of irreducible connections on the Dynkin diagrams*, Expo. Math. **28** (2010), no. 3, 218–253, MR2670999 DOI:10.1016/j.exmath.2009.11.001.
- [GS12] Pinhas Grossman and Noah Snyder, *Quantum subgroups of the Haagerup fusion categories*, Comm. Math. Phys. **311** (2012), no. 3, 617–643, MR2909758, DOI:10.1007/s00220-012-1427-x.
- [Gup08] Ved Prakash Gupta, *Planar algebra of the subgroup-subfactor*, Proc. Indian Acad. Sci. Math. Sci. **118** (2008), no. 4, 583–612, MR2511128 DOI:10.1007/s12044-008-0046-0 arXiv:0806.1791.
- [Han10] Richard Han, *A Construction of the “2221” Planar Algebra*, ProQuest LLC, Ann Arbor, MI, 2010, Thesis (Ph.D.)—University of California, Riverside, MR2822034 arXiv:1102.2052.
- [HP17] André Henriques and David Penneys, *Bicommutant categories from fusion categories*, Selecta Math. (N.S.) **23** (2017), no. 3, 1669–1708, MR3663592 DOI:10.1007/s00029-016-0251-0 arXiv:1511.05226.
- [HPT16a] André Henriques, David Penneys, and James Tener, *Categorified trace for module tensor categories over braided tensor categories*, Doc. Math. **21** (2016), 1089–1149, MR3578212 arXiv:1509.02937.
- [HPT16b] André Henriques, David Penneys, and James E. Tener, *Planar algebras in braided tensor categories*, 2016, arXiv:1607.06041.
- [JMS14] Vaughan F. R. Jones, Scott Morrison, and Noah Snyder, *The classification of subfactors of index at most 5*, Bull. Amer. Math. Soc. (N.S.) **51** (2014), no. 2, 277–327, MR3166042, arXiv:1304.6141, DOI:10.1090/S0273-0979-2013-01442-3.
- [Jon83] Vaughan F. R. Jones, *Index for subfactors*, Invent. Math. **72** (1983), no. 1, 1–25, MR696688, DOI:10.1007/BF01389127.
- [Jon99] ———, *Planar algebras I*, 1999, arXiv:math.QA/9909027.
- [Jon00] ———, *The planar algebra of a bipartite graph*, Knots in Hellas ’98 (Delphi), Ser. Knots Everything, vol. 24, World Sci. Publ., River Edge, NJ, 2000, MR1865703, pp. 94–117.
- [Jon01] ———, *The annular structure of subfactors*, Essays on geometry and related topics, Vol. 1, 2, Monogr. Enseign. Math., vol. 38, Enseignement Math., Geneva, 2001, MR1929335, pp. 401–463.
- [Jon11] ———, *Jones’ notes on planar algebras*, <http://math.berkeley.edu/~vfr/VANDERBILT/p121.pdf>, 2011.
- [Jon12] ———, *Quadratic tangles in planar algebras*, Duke Math. J. **161** (2012), no. 12, 2257–2295, MR2972458, arXiv:1007.1158, DOI:10.1215/00127094-1723608.
- [JP11] Vaughan F. R. Jones and David Penneys, *The embedding theorem for finite depth subfactor planar algebras*, Quantum Topol. **2** (2011), no. 3, 301–337, arXiv:1007.3173, MR2812459, DOI:10.4171/QT/23.
- [JP19] Corey Jones and David Penneys, *Realizations of algebra objects and discrete subfactors*, Adv. Math. **350** (2019), 588–661, MR3948170 DOI:10.1016/j.aim.2019.04.039 arXiv:1704.02035.
- [LMP15] Zhengwei Liu, Scott Morrison, and David Penneys, *1-Supertransitive Subfactors with Index at Most $6\frac{1}{5}$* , Comm. Math. Phys. **334** (2015), no. 2, 889–922, MR3306607, arXiv:1310.8566, DOI:10.1007/s00220-014-2160-4.
- [LR97] R. Longo and J. E. Roberts, *A theory of dimension*, K-Theory **11** (1997), no. 2, 103–159, MR1444286 DOI:10.1023/A:1007714415067.
- [MP14] Scott Morrison and Emily Peters, *The little desert? Some subfactors with index in the interval $(5, 3 + \sqrt{5})$* , Internat. J. Math. **25** (2014), no. 8, 1450080 (51 pages), MR3254427 DOI:10.1142/S0129167X14500803 arXiv:1205.2742.
- [MP15a] Scott Morrison and David Penneys, *2-supertransitive subfactors at index $3 + \sqrt{5}$* , J. Funct. Anal. **269** (2015), no. 9, 2845–2870, MR3394622 DOI:10.1016/j.jfa.2015.06.023 arXiv:1406.3401.
- [MP15b] ———, *Constructing spoke subfactors using the jellyfish algorithm*, Trans. Amer. Math. Soc. **367** (2015), no. 5, 3257–3298, MR3314808 DOI:10.1090/S0002-9947-2014-06109-6 arXiv:1208.3637.

- [MPS10] Scott Morrison, Emily Peters, and Noah Snyder, *Skein theory for the D_{2n} planar algebras*, J. Pure Appl. Algebra **214** (2010), no. 2, 117–139, [arXiv:0808.0764](#) [MR2559686](#) [DOI:10.1016/j.jpaa.2009.04.010](#).
- [MvN43] F. J. Murray and J. von Neumann, *On rings of operators. IV*, Ann. of Math. (2) **44** (1943), 716–808, [MR0009096](#).
- [nLa18] nLab authors, *principle of equivalence*, <http://ncatlab.org/nlab/show/principle%20of%20equivalence>, July 2018, [Revision 89](#).
- [NT13] Sergey Neshveyev and Lars Tuset, *Compact quantum groups and their representation categories*, Cours Spécialisés [Specialized Courses], vol. 20, Société Mathématique de France, Paris, 2013, [MR3204665](#).
- [Pen13] David Penneys, *A Planar Calculus for Infinite Index Subfactors*, Comm. Math. Phys. **319** (2013), no. 3, 595–648, [MR3040370](#) [arXiv:1110.3504](#) [DOI:10.1007/s00220-012-1627-4](#).
- [Pet10] Emily Peters, *A planar algebra construction of the Haagerup subfactor*, Internat. J. Math. **21** (2010), no. 8, 987–1045, [MR2679382](#), [DOI:10.1142/S0129167X10006380](#), [arXiv:0902.1294](#).
- [Pop94] Sorin Popa, *Classification of amenable subfactors of type II*, Acta Math. **172** (1994), no. 2, 163–255, [MR1278111](#), [DOI:10.1007/BF02392646](#).
- [PP15] David Penneys and Emily Peters, *Calculating two-strand jellyfish relations*, Pacific J. Math. **277** (2015), no. 2, 463–510, [MR3402358](#) [DOI:10.2140/pjm.2015.277-2](#) [arXiv:1308.5197](#).
- [Sel11] P. Selinger, *A survey of graphical languages for monoidal categories*, New structures for physics, Lecture Notes in Phys., vol. 813, Springer, Heidelberg, 2011, [MR2767048](#) [DOI:10.1007/978-3-642-12821-9_4](#), pp. 289–355.
- [Vic11] Jamie Vicary, *Categorical formulation of finite-dimensional quantum algebras*, Comm. Math. Phys. **304** (2011), no. 3, 765–796, [MR2794547](#) [DOI:10.1007/s00220-010-1138-0](#) [arXiv:0805.0432](#).
- [Yam04] Shigeru Yamagami, *Frobenius duality in C^* -tensor categories*, J. Operator Theory **52** (2004), no. 1, 3–20, [MR2091457](#).
- [Yam12] ———, *Representations of multicategories of planar diagrams and tensor categories*, 2012, [arXiv:1207.1923](#).